

ON ORDERS OF ELEMENTS OF FINITE ALMOST SIMPLE GROUPS WITH LINEAR OR UNITARY SOCLE

M.A. GRECHKOSEEVA

ABSTRACT. We say that a finite almost simple G with socle S is *admissible* (with respect to the spectrum) if G and S have the same sets of orders of elements. Let L be a finite simple linear or unitary group of dimension at least three over a field of odd characteristic. We describe admissible almost simple groups with socle L . Also we calculate the orders of elements of the coset $L\tau$, where τ is the inverse-transpose automorphism of L .

1. INTRODUCTION

The *spectrum* $\omega(G)$ of a finite group G is the set of orders of its elements, and groups with equal spectra are said to be *isospectral*. This paper is a part of a larger investigation devoted to recognition of simple groups by spectrum. To solve the *problem of recognition by spectrum* for a given finite nonabelian simple group S is to describe (up to isomorphism) finite groups that are isospectral to S . As a working hypothesis of that investigation, it was conjectured that a finite group G isospectral to a “sufficiently large” simple group S must be an almost simple group with socle S , i. e. $S \leq G \leq \text{Aut } S$. In 2015, this conjecture was proved with the following precise meaning of the term “sufficiently large”: S is a linear or unitary group of dimension larger than 44, or S is a symplectic or orthogonal group of dimension larger than 60, or S is one of the sporadic, alternating and exceptional groups of Lie type other than J_2 , A_6 , A_{10} , and ${}^3D_4(2)$ (see [13]). Thus for a vast class of simple groups, the initial problem of recognition by spectrum was reduced to a more specific problem of describing (up to isomorphism) almost simple groups with socle S that are isospectral to S , and this is the problem we address in this paper.

For brevity, we refer to a finite almost simple group with socle S that is isospectral to S as *admissible* for S . Clearly we are interested in non-trivial admissible groups, i. e. other than S itself. It is not hard to check that there are no non-trivial admissible groups for the alternating groups. The information collected in [6] allows to verify that the sporadic groups do not possess non-trivial admissible groups either. One of the first examples of non-trivial admissible groups was discovered by Mazurov [15]: he showed that the finite groups isospectral to $PSL_3(5)$ are exactly $PSL_3(5)$ and its extension by the graph automorphism. Later Zavarnitsine [19] provided an example demonstrating that the number of admissible groups can be arbitrarily large: $PSL_3(7^{3^k})$ has exactly $k + 1$ admissible groups, including itself, and these are precisely extensions by field automorphisms.

Admissible groups are described for all exceptional groups of Lie type (see [22] for references), $PSL_2(q)$ (see [1]), $PSL_3(q)$ and $PSU_3(q)$ (see [19, 20]), classical groups over fields of characteristic 2 [10, 12, 23], and symplectic and odd-dimensional orthogonal groups

2010 *Mathematics Subject Classification.* 20D06, 20D60.

Key words and phrases. Almost simple group, linear group, unitary group, orders of elements, inverse-transpose map.

The work is supported by Russian Science Foundation (project 14-21-00065).

over fields of odd characteristic [11]. It is worth noting that for all of these groups, there is $\alpha \in \text{Aut } S$ such that $S_0 = \langle S, \alpha \rangle$ is admissible and any other admissible group is conjugate in $\text{Aut } S$ to a subgroup of S_0 ; in other words, any admissible group is a cyclic extension of S and up to isomorphism there is a unique maximal admissible group. Below we will see that not all simple groups satisfy the latter property.

The main result of this paper is a description of admissible groups for linear and unitary groups over fields of odd characteristic (Theorems 1, 2 and 3). Also we explicitly describe spectra of some almost simple groups with linear or unitary socle (Lemmas 3.3, 4.6, and 4.7). In the rest of this section, we introduce the notation used in the theorems and then state our results.

Throughout this paper, p is a prime, F is the algebraic closure of the field of order p , $H = GL_n(F)$, with matrices acting on row vectors by right multiplication, and τ is the inverse-transpose map $g \mapsto g^{-\top}$ of H . If q is a power of p , then F_q denotes the subfield of F of order q , λ_q denotes a fixed primitive element of F_q and φ_q denotes the standard Frobenius endomorphism of H of level q , i. e. the endomorphism induced by raising matrix entries to the q th power. We identify $GL_n(q)$ with $C_H(\varphi_q)$ and $GU_n(q)$ with $C_H(\varphi_q\tau)$.

We write $GL_n^+(q)$ instead of $GL_n(q)$ and $GL_n^-(q)$ instead of $GU_n(q)$ and use a similar agreement for $PGL_n(q)$, $SL_n(q)$, and $PSL_n(q)$. For $\varepsilon \in \{+, -\}$, we shorten $\varepsilon 1$ to ε in arithmetic expressions.

As usual, by $(a_1, \dots, a_s)$ we denote the greatest common divisor of positive integers $a_1, \dots, a_s$, and by $[a_1, \dots, a_s]$ we denote their least common multiple. If a and b are positive integers, then $\pi(a)$ denotes the set of prime divisors of a , $(a)_b$ denotes the largest divisor c of a such that $\pi(c) \subseteq \pi(b)$ and $(a)_{b'}$ denotes the number $a/(a)_b$.

Let $L = PSL_n^\varepsilon(q)$, where $q = p^m$, and define $d = (n, q - \varepsilon)$. We write $\delta = \delta(\varepsilon q)$ to denote the diagonal automorphism of L induced by $\text{diag}(\lambda, 1, \dots, 1)$, where λ is a primitive $q - \varepsilon$ th root of unity in F_{q^2} . We denote by φ the field automorphism induced by φ_p . The automorphism induced by τ is denoted by the same letter. The image of $\alpha \in \text{Aut } L$ in $\text{Out } L$ is denoted by $\bar{\alpha}$.

When $n \geq 3$, the inverse-transpose automorphism is outer and $\text{Out } L$ has the following presentation (we omit overbars for convenience):

$$\begin{aligned} \langle \delta, \varphi, \tau \mid \delta^d = \varphi^m = \tau^2 = [\varphi, \tau] = 1, \delta^\varphi = \delta^p, \delta^\tau = \delta^{-1} \rangle & \quad \text{if } \varepsilon = +, \\ \langle \delta, \varphi, \tau \mid \delta^d = \tau^2 = [\varphi, \tau] = 1, \varphi^m = \tau, \delta^\varphi = \delta^p, \delta^\tau = \delta^{-1} \rangle & \quad \text{if } \varepsilon = -. \end{aligned}$$

Theorem 1 is concerned with the extension by the inverse-transpose automorphism τ . A criterion of admissibility of this extension is not very short, so it seems reasonable to write up it separately.

Theorem 1. *Let $L = PSL_n^\varepsilon(q)$, where $n \geq 3$, $\varepsilon \in \{+, -\}$ and q is odd, and $G = L \rtimes \langle \tau \rangle$. Then either $\omega(G) = \omega(L)$ or one of the following holds:*

- (i) $n = p^{t-1} + 2$ with $t \geq 1$, $q \equiv -\varepsilon \pmod{4}$ and $4p^t \in \omega(G) \setminus \omega(L)$;
- (ii) $n = 2^t + 1$ with $t \geq 1$, $(n, q - \varepsilon) > 1$ and $2(q^{(n-1)/2} - \varepsilon^{(n-1)/2}) \in \omega(G) \setminus \omega(L)$;
- (iii) $n = p^{t-1} + 1$ with $t \geq 1$ and $2p^t \in \omega(G) \setminus \omega(L)$;
- (iv) n is even, $(n)_2 \leq (q - \varepsilon)_2$, $q \equiv \varepsilon \pmod{4}$ and $q^{n/2} + \varepsilon^{n/2} \in \omega(G) \setminus \omega(L)$;
- (v) n is even, $(n)_{2'} > 3$, $(n, q - \varepsilon)_{2'} > 1$ and $2[q^{(n)_2} - 1, q^{n/2-(n)_2} + \varepsilon^{n/2-(n)_2}] \in \omega(G) \setminus \omega(L)$.

TABLE 1.

Conditions on L		$\alpha = \gamma\beta = \beta\gamma$	
		γ	β
$n = p^t + 1,$	$(b)_2 > 2$ or $(n)_2 < (p-1)_2$	1	$\varphi^{m/2}\tau\eta$
$n \neq 2^u + 2$	$(m)_2 = 2, (n)_2 < (p+1)_2$	1	$\varphi^{m/2}\eta$
$n \neq p^t + 1,$	τ not admissible, $ \psi > 1$	ψ	1
b odd	τ admissible	ψ	τ
$n \neq p^t + 1,$ b even, $p \equiv \epsilon \pmod{4}$	$n = p^s + 2^u + 1$	ψ	$\phi, \phi^i\tau$
	$n \neq p^s + 2^u + 1, (n)_2 \geq (p-\epsilon)_2$	ψ	$\phi, \phi^i\tau, \phi^{2j}\tau\eta$
	$n \neq p^s + 2^u + 1, (n)_2 < (p-\epsilon)_2, \epsilon = +$	ψ	$\phi, \phi^i\tau, \phi^{2j}\tau\eta, \phi\tau\eta$
	$n \neq p^s + 2^u + 1, (n)_2 < (p-\epsilon)_2, \epsilon = -$	ψ	$\phi, \phi^i\tau, \phi^{2j}\tau\eta, \phi\eta$
$t, u > 0, s \geq 0$		$2i \mid (b)_2, 4j \mid (b)_2$	

Theorems 2 and 3 describe admissible groups appealing to Theorem 1. As we mentioned, the admissible groups for $PSL_3(q)$ and $PSU_3(q)$ were determined by Zavarnitsine [19, 20], so we do not consider these groups. However, we include them into the statements of the theorems for completeness. Observe that two almost simple groups with socle S are isomorphic if and only if their images in $\text{Out } S$ are conjugate. Thus to describe admissible groups up to isomorphism, it is sufficient to describe their images in $\text{Out } S$ up to conjugacy. We refer to $\alpha \in \text{Aut } S$ as admissible if $\langle S, \alpha \rangle$ is admissible.

Theorem 2. *Let $L = PSL_n(q)$, where $n \geq 3$, $q = p^m$ and p is an odd prime. Let $d = (n, q-1)$, $b = ((q-1)/d, m)_d$, $\eta = \delta^{(d)_{2'}}$, $\phi = \varphi^{m/(b)_2}$, and*

$$\psi = \begin{cases} \varphi^{m/(b)_{2'}} & \text{if } n \neq 4 \text{ or } 12 \nmid q+1; \\ \varphi^{(m)_{3'}} & \text{if } n = 4 \text{ or } 12 \mid q+1. \end{cases}$$

Suppose that $L < G \leq \text{Aut } L$. Then $\omega(G) = \omega(L)$ if and only if G/L is conjugate in $\text{Out } L$ to a subgroup of $\langle \bar{\alpha} \rangle$, where α is one of the elements specified in Table 1.

As an example of applying Theorem 2, let us consider admissible groups for $PSL_4(25)$. For this group, $d = 4$, $b = 2$, $\phi = \varphi$, $n \neq p^t + 1$ and $n = p^0 + 2^1 + 1$. Thus up to isomorphism there are two non-trivial admissible groups, namely, the extensions by φ and by $\varphi\tau$. Clearly they are both maximal.

Theorem 3. *Let $L = PSU_n(q)$, where $n \geq 3$, $q = p^m$ and p be an odd prime. Let $d = (n, q+1)$, $b = ((q+1)/d, m)_d$ and*

$$\psi = \begin{cases} \varphi^{2m/(b)_{2'}} & \text{if } n \neq 4 \text{ or } 12 \nmid q-1; \\ \varphi^{2(m)_{3'}} & \text{if } n = 4 \text{ and } 12 \mid q-1. \end{cases}$$

Suppose that $L < G \leq \text{Aut } L$. Then $\omega(G) = \omega(L)$ if and only if $n-1$ is not a power of p and G/L is conjugate in $\text{Out } L$ to a subgroup of $\langle \bar{\alpha} \rangle$, where

- (i) $\alpha = \psi$ if τ is not admissible and $|\psi| > 1$;
- (ii) $\alpha = \psi\tau$ if τ is admissible, $(n)_2 > 2$ and $n \geq 16$;
- (iii) $\alpha = \psi\varphi^{(m)_{2'}}$ if τ is admissible and either $(n)_2 \leq 2$ or $n \leq 12$.

Returning now to the initial recognition problem, we state the following consequence of [17, Theorem 1] and the above results.

Corollary. *Let $L = PSL_n^\varepsilon(q)$, where $n \geq 45$, $\varepsilon \in \{+, -\}$ and q is odd. A finite group is isospectral to L if and only if it is isomorphic to an almost simple group G with socle L and $G/L = \langle \bar{\alpha} \rangle$, where α is an identity or is as specified in Theorems 2 and 3.*

2. SPECTRA OF CLASSICAL GROUPS AND RELATED NUMBER-THEORETICAL LEMMAS

In this section, we collect necessary information on spectra of classical groups and related number-theoretical lemmas. Our notation for the classical groups follows that of [6]. Recall some well-known isomorphisms between classical groups (see, for example, [14, Proposition 2.9.1]). If q is odd, then

$$\begin{aligned} PSp_4(q) &\simeq \Omega_5(q), \quad \Omega_6^\varepsilon(q) \simeq SL_4^\varepsilon(q)/\{\pm 1\}, \quad P\Omega_6^\varepsilon(q) \simeq PSL_4^\varepsilon(q), \\ \Omega_4^+(q) &\simeq SL_2(q) \circ SL_2(q), \quad P\Omega_4^+(q) \simeq PSL_2(q) \times PSL_2(q), \quad \Omega_4^-(q) \simeq PSL_2(q^2), \\ Sp_2(q) &\simeq SL_2(q), \quad \Omega_3(q) \simeq PSL_2(q). \end{aligned} \quad (2.1)$$

Given a prime r , we write $\omega_{r'}(G)$ to denote the set of orders of elements of G that are coprime to r . In particular, if G is a group of Lie type over a field of characteristic p , then $\omega_p(G)$ is the set of orders of semisimple elements of G . By $\omega_{\bar{r}}(G)$ we denote the difference $\omega(G) \setminus \omega_{r'}(G)$.

Lemma 2.1. *Let $n \geq 2$, q be a power of a prime p , $\varepsilon \in \{+, -\}$ and let G be $PGL_n^\varepsilon(q)$ or $PSL_n^\varepsilon(q)$. Let $d = 1$ in the first case and $d = (n, q - 1)$ in the second case. Then $\omega(G)$ consists of all divisors of the following numbers:*

- (i) $(q^n - \varepsilon^n)/((q - \varepsilon)d)$;
- (ii) $[q^{n_1} - \varepsilon^{n_1}, q^{n_2} - \varepsilon^{n_2}]/(n/(n_1, n_2), d)$, where $n_i > 0$ and $n_1 + n_2 = n$;
- (iii) $[q^{n_1} - \varepsilon^{n_1}, \dots, q^{n_s} - \varepsilon^{n_s}]$, where $s \geq 2$, $n_i > 0$ and $n_1 + \dots + n_s = n$;
- (iv) $p^t(q^{n_1} - \varepsilon^{n_1})/d$, where $t \geq 1$, $n_1 > 0$ and $p^{t-1} + 1 + n_1 = n$;
- (v) $p^t[q^{n_1} - \varepsilon^{n_1}, \dots, q^{n_s} - \varepsilon^{n_s}]$, where $t \geq 1$, $s \geq 2$, $n_i > 0$ and $p^{t-1} + 1 + n_1 + \dots + n_s = n$;
- (vi) $(n, q - 1)p^t/d$ if $n = p^{t-1} + 1$ for some $t \geq 1$.

Proof. See [3, Corollaries 2 and 3]. □

Lemma 2.2. *Let $n \geq 1$, q be a power of an odd prime p and let G be one of the groups $Sp_{2n}(q)$, $PSp_{2n}(q)$, and $\Omega_{2n+1}(q)$. Let $d = c = 1$ if $G = Sp_{2n}(q)$; $d = 2$ and $c = 1$ if $G = PSp_{2n}(q)$ or $G = \Omega_5(q), \Omega_3(q)$; and $d = c = 2$ if $G = \Omega_{2n+1}(q)$ and $n \geq 3$. Then $\omega(G)$ consists of all divisors of the following numbers:*

- (i) $(q^n \pm 1)/d$;
- (ii) $[q^{n_1} \pm 1, \dots, q^{n_s} \pm 1]$, where $s \geq 2$, $n_i > 0$ and $n_1 + \dots + n_s = n$;
- (iii) $p^t(q^{n_1} \pm 1)/c$, where $t \geq 1$, $n_1 > 0$ and $p^{t-1} + 1 + 2n_1 = 2n$;
- (iv) $p^t[q^{n_1} \pm 1, \dots, q^{n_s} \pm 1]$, where $t \geq 1$, $s \geq 2$, $n_i > 0$ and $p^{t-1} + 1 + 2n_1 + \dots + 2n_s = 2n$;
- (v) $2p^t/d$ if $2n = p^{t-1} + 1$ for some $t \geq 1$.

Proof. See [4, Corollaries 1, 2 and 6] for $n \geq 2$ and (2.1) together with Lemma 2.1 for $n = 1$. □

Lemma 2.3. *Let $n \geq 2$, q be a power of an odd prime p and $\varepsilon \in \{+, -\}$. Then $\omega_{p'}(\Omega_{2n}^\varepsilon(q))$ consists of all divisors of the following numbers:*

- (i) $(q^n - \varepsilon)/2$;
- (ii) $[q^{n_1} - \kappa_1, \dots, q^{n_s} - \kappa_s]$, where $s \geq 2$, $\kappa_i \in \{+, -\}$, $n_i > 0$, $n_1 + \dots + n_s = n$ and $\kappa_1 \kappa_2 \dots \kappa_s = \varepsilon$;

and $\omega_{p'}(P\Omega_{2n}^\varepsilon(q))$ consists of all divisors of the following numbers:

- (i) $(q^n - \varepsilon)/(4, q^n - \varepsilon)$;
- (ii) $[q^{n_1} - \kappa, q^{n_2} - \varepsilon \kappa]/e$, where $\kappa \in \{+, -\}$, $n_1, n_2 > 0$, $n_1 + n_2 = n$; $e = 2$ when $(q^{n_1} - \kappa)_2 = (q^{n_2} - \varepsilon \kappa)_2$ and $e = 1$ otherwise;
- (iii) $[q^{n_1} - \kappa_1, \dots, q^{n_s} - \kappa_s]$, where $s \geq 3$, $\kappa_i \in \{+, -\}$, $n_i > 0$, $n_1 + \dots + n_s = n$ and $\kappa_1 \kappa_2 \dots \kappa_s = \varepsilon$.

Proof. For $n \geq 4$, see [5, Theorem 6]. If $n = 2, 3$, the assertion follows from the isomorphisms given in (2.1) and Lemma 2.1. $\square$

Let $k \geq 3$ and q be an integer whose absolute value is greater than one. A primitive prime divisor of $q^k - 1$ is a prime r such that r divides $q^k - 1$ and does not divide $q^i - 1$ for any $i < k$. The set of primitive prime divisors of $q^k - 1$ is denoted by $R_k(q)$, and $r_k(q)$ denotes some fixed element of $R_k(q)$.

Lemma 2.4 (Zsigmondy [21]). *Let q be a prime power, $\varepsilon \in \{+, -\}$, and $k \geq 3$. Suppose that $(q, \varepsilon, k) \neq (2, +, 6), (2, -, 3)$. Then $R_k(q)$ is not empty.*

Two following results are well-known.

Lemma 2.5. *Let $q > 1$, k and l be positive integers, and $\varepsilon \in \{+, -\}$. Then*

- (i) $(q^k - 1, q^l - 1) = q^{(k,l)} - 1$;
- (ii) $(q^k + 1, q^l + 1)$ is equal to $q^{(k,l)} + 1$ if $(k)_2 = (l)_2$ and to $(2, q + 1)$ otherwise;
- (iii) $(q^k - 1, q^l + 1)$ is equal to $q^{(k,l)} + 1$ if $(k)_2 > (l)_2$ and to $(2, q + 1)$ otherwise;
- (iv) $((q^k - \varepsilon^k)/(q - \varepsilon), k) = (q - \varepsilon, k)$;
- (v) if $(k, l) = 1$, then $(q^l - \varepsilon^l)/(q - \varepsilon)$ divides $(q^{lk} - \varepsilon^{lk})/(q^k - \varepsilon^k)$ and $(q^l - \varepsilon^l)/(n, q - \varepsilon)$ divides $(q^{lk} - \varepsilon^{lk})/(n, q^k - \varepsilon^k)$ for any positive integer n .

Lemma 2.6. *Let $q > 1$ and k be positive integers and $\varepsilon \in \{+, -\}$.*

- (i) *If an odd prime r divides $q - \varepsilon$, then $(q^k - \varepsilon^k)_r = (k)_r(q - \varepsilon)_r$.*
- (ii) *If an odd prime r divides $q^k - \varepsilon^k$, then r divides $q^{(k)_{r'}} - \varepsilon^{(k)_{r'}}$.*
- (iii) *If 4 divides $q - \varepsilon$ and k is odd, then $(q^k - \varepsilon^k)_2 = (k)_2(q - \varepsilon)_2$.*

Lemma 2.7. *Let q be odd and $n \geq 4$ be even. Then $q^{n/2} + \varepsilon^{n/2} \in \omega(PSL_n^\varepsilon(q))$ if and only if $(n)_2 > (q - \varepsilon)_2$.*

Proof. Let $d = (n, q - \varepsilon)$. Since $a = q^{n/2} + \varepsilon^{n/2}$ is divisible by a primitive divisor $r_n(\varepsilon q)$, it follows from 2.1 that $a \in \omega(PSL_n^\varepsilon(q))$ if and only if a divides

$$\frac{q^n - 1}{(q - \varepsilon)d} = \frac{(q^{n/2} + \varepsilon^{n/2})(q^{n/2} - \varepsilon^{n/2})}{(q - \varepsilon)d};$$

that is, if and only if d divides $(q^{n/2} - \varepsilon^{n/2})/(q - \varepsilon)$.

If $n/2$ is odd, then by Lemma 2.6, we have that $(q^{n/2} - \varepsilon^{n/2})/(q - \varepsilon)$ is odd and, therefore, it is not divisible by d , which is even. If $n/2$ is even and $(n)_2 \leq (q - \varepsilon)_2$, then

$q \equiv \varepsilon \pmod{4}$ and $(d)_2 = (n)_2$, and hence $(q^{n/2} - \varepsilon^{n/2})_2 / (q - \varepsilon)_2 = (n)_2 / 2 < (d)_2$. Also, if $(n)_2 > (q - \varepsilon)_2$, then

$$((q^{n/2} - \varepsilon) / (q - \varepsilon), d) = ((q^{n/2} - \varepsilon) / (q - \varepsilon), q - \varepsilon, n) = (n/2, q - \varepsilon, n) = d,$$

where the second equality holds by Lemma 2.5(iv). $\square$

3. EXTENSIONS BY FIELD AND GRAPH-FIELD AUTOMORPHISMS

In this section, we derive some formulas concerning orders of elements in extensions of $PSL_n^\varepsilon(q)$ by field or graph-field automorphisms. Following the lines of the proof of [20, Proposition 13], we will exploit a correspondence between σ -conjugacy classes of $C_K(\sigma^k)$ and conjugacy classes of $C_K(\sigma)$, where K is a connected linear algebraic group and σ is a Steinberg endomorphism of K , i.e. a surjective endomorphism with finitely many fixed points. Also we will use a slight modification of this correspondence inspired by [9, Theorem 2.1].

We begin with necessary notation and the Lang–Steinberg theorem. If K is a group and σ is an endomorphism of K , then we write K_σ to denote $C_K(\sigma)$.

Lemma 3.1 (Lang–Steinberg). *Let K be a connected linear algebraic group and σ be a surjective endomorphism of K such that K_σ is finite. Then the map $x \mapsto x^{-\sigma}x$ from K to K is surjective.*

Recall that F is the algebraic closure of the field of order p and φ_q is the endomorphism of $GL_n(F)$ raising matrix entries to the q th power, where q is a power of p . An endomorphism σ of a linear algebraic group K is said to be a Frobenius endomorphism if there are an identification of K with a closed subgroup of $GL_n(F)$ and a positive integer k such that σ^k is induced by φ_q . Clearly, if σ is a Frobenius endomorphism, then σ is a Steinberg endomorphism.

Lemma 3.2. *Let K be a connected linear algebraic group over F , α be a Frobenius endomorphism of K and τ be an automorphism of K of order 2 that commutes with α . Let k be a positive integer, $l \in \{0, 1\}$ and let β be the automorphism of $K_{\alpha^k \tau^l}$ induced by α . Given $g \in K_{\alpha^k \tau^l}$, choose $z \in K$ such that $g = z^{-\alpha}z$ and define $\zeta(g) = z^{\tau^l} z^{-\alpha^k}$. Then $\zeta(g) \in K_\alpha$, $(\beta g)^k = z^{-1} \tau^l \zeta(g) z$ and the map $[\beta g] \mapsto [\tau^l \zeta(g)]$ is a one-to-one correspondence between the $K_{\alpha^k \tau^l}$ -conjugacy classes in the coset $\beta K_{\alpha^k \tau^l}$ and the K_α -conjugacy classes in the coset $\tau^l K_\alpha$.*

Proof. Observe that α^k and $\alpha^k \tau$ are Frobenius endomorphisms of K .

Let $g, z \in K$, $g = z^{-\alpha}z$ and $h = z^{\tau^l} z^{-\alpha^k}$. Define $f = \tau^l h = z^{\tau^l} z^{-\alpha^k}$. Then

$$g^{\alpha^k \tau^l} = g \Leftrightarrow \tau^l z^{-\alpha^{k+1}} z^{\alpha^k} \tau^l = z^{-\alpha} z \Leftrightarrow z^{\alpha} \tau^l z^{-\alpha^{k+1}} = z^{\tau^l} z^{-\alpha^k} \Leftrightarrow f^\alpha = f. \quad (3.1)$$

This if $g \in K_{\alpha^k \tau^l}$, then $f \in \tau^l K_\alpha$. Furthermore, $(\beta g)^k = \beta^k g^{\alpha^{k-1}} \dots g = \tau^l z^{-\alpha^k} z = z^{-1} f z$.

Let $g_1 = z_1^{-\alpha} z_1 \in K_{\alpha^k \tau^l}$ and $g_2 = z_2^{-\alpha} z_2 \in K_{\alpha^k \tau^l}$. Suppose that $x^{-1} \alpha g_1 x = \alpha g_2$ for some $x \in K_{\alpha^k \tau^l}$. Then writing $y = z_1 x z_2^{-1}$, we have $y^\alpha = y$ and

$$f_1 = z_1 (\beta g_1)^k z_1^{-1} = z_1 x (\beta g_2)^k x^{-1} z_1^{-1} = y f_2 y^{-1}.$$

Conversely, if $f_1 = y f_2 y^{-1}$ with $y \in K_\alpha$, then $x = z_1^{-1} y z_2$ satisfies $x^{\alpha^k \tau^l} = x$ and

$$x^{-1} \alpha g_1 x = z_2^{-1} y^{-1} z_1 \alpha z_1^{-\alpha} z_1^{-1} y z_2 = z_2^{-1} y^{-1} \alpha y z_2 = z_2^{-1} \alpha z_2 = \alpha g_2.$$

It follows that the map under consideration translate conjugacy classes to conjugacy classes and is injective. Furthermore, the conjugacy class $[f]$ does not depend on the choice of z in the equality $g = z^{-\alpha}z$.

Now let $h \in K_\alpha$. By the Lang–Steinberg theorem, there is $z \in K$ such that $h^{-\tau^l} = z^{\alpha^k \tau^l} z^{-1}$, and therefore $\tau^l h = z \tau^l z^{-\alpha^k}$. Then (3.1) implies that $g = z^{-\alpha}z$ lies in $K_{\alpha^k \tau^l}$, with $[\beta g]$ mapping to $[\tau^l h]$. Thus the map is surjective, and the proof is complete. $\square$

It should be noted that some special cases of Lemma 3.2 were proved in [7, Lemma 2.10] ($l = 0$) and [9, Theorem 2.1] ($l = 1$ and $k = 1$).

We use Lemma 3.2 to establish the following result generalizing [20, Corollary 14].

Lemma 3.3. *Let $L = PSL_n(q)$ and $U = PSU_n(q)$, where $n \geq 3$ and $q = p^m$. Let k divides m , $q = q_0^k$ and $\beta = \varphi^{m/k}$. Then for any i , we have the following:*

- (i) $\omega(\beta \delta^i(q)L) = k \cdot \omega(\delta^i(q_0)PSL_n(q_0))$;
- (ii) if k is even, then $\omega(\beta \tau \delta^i(q)L) = k \cdot \omega(\delta^i(-q_0)PSU_n(q_0))$;
- (iii) if k is odd, then $\omega(\beta \tau \delta^i(-q)U) = k \cdot \omega(\delta^i(-q_0)PSU_n(q_0))$;
- (iv) if k is odd, then $\omega(\beta \tau \delta^i(q)L) = k \cdot \omega(\tau \delta^i(q_0)PSL_n(q_0))$;
- (v) $\omega(\beta \delta^i(-q)U) = k \cdot \omega(\tau \delta^i(q_0)PSL_n(q_0))$.

In particular, $\omega(\tau \delta^i(q)L) = \omega(\tau \delta^i(-q)U)$.

Proof. Denote the center of $H = GL_n(F)$ by Z . In particular, we write $|gZ|$ to denote the projective order of a matrix $g \in H$.

We apply Lemma 3.2 to H with $\alpha = \varphi_p^{m/k}$ and $l = 0$. Observe that $H_{\alpha^k} = GL_n(q)$ and $H_\alpha = GL_n(q_0)$. Let $g \in GL_n(q)$ and $h = \zeta(g)$ be the element of $GL_n(q_0)$ defined in Lemma 3.2. Since $(\beta g)^k$ is conjugate to h in H , we see that

$$|\beta gZ| = k \cdot |(\beta gZ)^k| = k \cdot |hZ|,$$

and also

$$\det h = \det(g^{\beta^{k-1}} \dots g) = (\det g)^{q_0^{k-1} + \dots + 1} = (\det g)^{(q-1)/(q_0-1)}.$$

Let $\lambda = \lambda_q$ and $\lambda_0 = \lambda_{q_0}$. The condition $gZ \in \delta^i L$ is equivalent to $\det g \in \langle \lambda^i, \lambda^n \rangle = \langle \lambda^{(i,n)} \rangle$. Since $\lambda^{(q-1)/(q_0-1)}$ is a primitive element of F_{q_0} , it follows that $\det g \in \langle \lambda^{(i,n)} \rangle$ if and only if $\det h \in \langle \lambda_0^{(i,n)} \rangle$. Thus (i) holds. Similarly, (ii) and (iii) follow from Lemma 3.2 with $\alpha = \varphi_p^{m/k} \tau$, $l = 0$.

Let k be odd and $\alpha = \varphi_p^{m/k} \tau$. Then $H_{\alpha^k \tau} = GL_n(q)$ and $H_\alpha = GU_n(q_0)$. Applying Lemma 3.2 with $l = 1$, we construct from an element $g \in GL_n(q)$ an element $h \in GU_n(q_0)$ such that $g = z^{-\alpha}z$ and $h = z^\tau z^{-\alpha^k}$ for some $z \in H$. As in the previous case, we deduce that $|\beta \tau gZ| = k \cdot |\tau hZ|$. Furthermore, $\det g = (\det z)^{q_0+1}$ and $\det h = (\det z)^{q-1}$. Since $(\det g)^{q-1} = 1$, it follows that $(\det z)^{(q_0+1)(q-1)} = 1$. Let $\lambda = \lambda_q$, $\lambda_0 = \lambda_{q_0}^{q_0-1}$ and μ be a primitive $(q_0+1)(q-1)$ th root of unity in F . Then $\det g \in \langle \lambda^{(i,n)} \rangle$ if and only if $\det z \in \langle \mu^{(i,n)} \rangle$, which is equivalent to $\det h \in \langle \lambda_0^{(i,n)} \rangle$. Hence

$$\omega(\beta \tau \delta^i(q)L) = k \cdot \omega(\tau \delta^i(-q_0)PSU_n(q_0)). \quad (3.2)$$

Similarly, taking $\alpha = \varphi_p^{m/k}$ and $l = 1$, we prove (v). Applying (v) with $k = 1$ and observing that in this case β acts on U in the same way as τ , we have

$$\omega(\tau \delta^i(q)L) = \omega(\tau \delta^i(-q)U).$$

Now this equality and (3.2) imply (iv). □

Note that for unitary groups, (iii) and (v) of Lemma 3.3 cover all possibilities for an element $\gamma \in \langle \varphi \rangle$: if $|\gamma| = 2k$, then $\langle \gamma \rangle = \langle \varphi^{m/k} \rangle$; while if $|\gamma| = k$ with k odd, then $\langle \gamma \rangle = \langle \varphi^{m/k} \tau \rangle$ because $(m + m/k, 2m) = 2m/k$. Thus Lemma 3.3 expresses the spectrum of an extension by a field or graph-field automorphism in terms of known spectra and the spectrum of the extension by graph automorphism, which we consider in the next section.

4. EXTENSION BY GRAPH AUTOMORPHISM

This section is largely concerned with matrices, so we need to define some of them. We denote by E the identity matrix whose size is clear from the context, and by J_k the $k \times k$ unipotent Jordan block.

Recall that $\langle \bar{\delta} \rangle \rtimes \langle \bar{\tau} \rangle$ is a dihedral group of order $2(n, q - \varepsilon)$. It follows that for odd n , every $\tau \delta^i$ is conjugate to τ modulo L , and hence

$$\omega(\tau L) = \omega(\cup_{i=1}^d \tau \delta^i L) = \omega(\tau PGL_n(q)). \quad (4.1)$$

If n is even, then for any i , we have

$$\omega(\tau L) = \omega(\tau \delta^{2i} L) \quad \text{and} \quad \omega(\tau \delta L) = \omega(\tau \delta^{2i-1} L). \quad (4.2)$$

As we saw in Lemma 3.3, the cosets $\tau \delta^i(q)PSL_n(q)$ and $\tau \delta^i(-q)PSU_n(q)$ have the same orders of elements, so it suffices to describe $\omega(\tau \delta^i(q)PSL_n(q))$.

Since $|g\tau| = 2|(g\tau)^2| = 2|gg^\tau|$, it follows that the elements of $\omega(\tau PGL_n(q))$ are exactly twice the projective orders of elements of

$$\Gamma_n(q) = \{gg^\tau \mid g \in GL_n(q)\}.$$

If n is even, then $\omega(\tau L)$ and $\omega(\tau \delta L)$ are analogously related to the projective orders of elements of

$$\Gamma_n^\square(q) = \{gg^\tau \mid g \in GL_n(q), \det g \in (F_q^\times)^2\}$$

and

$$\Gamma_n^\boxtimes(q) = \{gg^\tau \mid g \in GL_n(q), \det g \notin (F_q^\times)^2\},$$

respectively.

A comprehensive treatment of the equation $h = gg^\tau$ for a given matrix h is provided by Fulman and Guralnick in [8], and we use the terminology and some results of this paper. First of all, it is helpful to note that $h = gg^\tau$ yields

$$xhx^{-1} = (xgx^\tau)(xgx^\tau)^\tau \quad (4.3)$$

for any $x \in GL_n(q)$, and hence h lies in $\Gamma_n(q)$, $\Gamma_n^\square(q)$ or $\Gamma_n^\boxtimes(q)$ if and only if the whole conjugacy class $[h]$ lies in the corresponding set. Thus we may work not with individual matrices but with conjugacy classes. Recall that the conjugacy classes of $GL_n(q)$ are parametrized by collections of partitions

$$\{\lambda_\phi \mid \phi \text{ is a monic irreducible polynomial over } F_q\}$$

such that $|\lambda_z| = 0$ and $\sum_\phi \deg(\phi)|\lambda_\phi| = n$. In this parametrization, the collection of partitions $\{\lambda_\phi\}$ corresponds to the class $[h]$ such that the multiplicity of ϕ^k as an elementary divisor of h is equal to the multiplicity of parts of size k in λ_ϕ . We denote the partition corresponding under this parametrization to a class $[h]$ and a polynomial ϕ by $\lambda_\phi(h)$.

A criterion for a matrix h to lie in $\Gamma_n(q)$ was obtained by Wall in [18]. In the same paper, Wall described the conjugacy classes of finite symplectic and orthogonal groups over fields of odd characteristic, and it turns out that the matrices of $\Gamma_n(q)$ are very similar to symplectic and orthogonal ones.

Lemma 4.1. *If q is odd, then $h \in \Gamma_n(q)$ if and only if h satisfies the following:*

- (i) h is conjugate to h^{-1} ;
- (ii) all even parts of $\lambda_{z-1}(h)$ have even multiplicity;
- (iii) all odd parts of $\lambda_{z+1}(h)$ have even multiplicity.

Proof. See [18, Theorem 2.3.1]. □

Lemma 4.2. *Let $h \in GL_n(q)$ and q be odd. Then h is conjugate to a element of $Sp_n(q)$ if and only if h satisfies the following:*

- (i) h is conjugate to h^{-1} ;
- (ii) all odd parts of $\lambda_{z-1}(h)$ and $\lambda_{z+1}(h)$ have even multiplicity.

Proof. See [18, p. 36, case (B)(i)]. □

Lemma 4.3. *Let $h \in GL_n(q)$ and q be odd. Then h is conjugate to an element of $GO_n^\varepsilon(q)$ for some ε if and only if h satisfies the following:*

- (i) h is conjugate to h^{-1} ;
- (ii) all even parts of $\lambda_{z-1}(h)$ and $\lambda_{z+1}(h)$ have even multiplicity.

Suppose that n is even and h is a unipotent element satisfying (ii). If $\lambda_{z-1}(h)$ has odd parts, then h is conjugate to an element of $GO_n^+(q)$ and also to an element of $GO_n^-(q)$. If there are no odd parts, then h is not conjugate to an element of $GO_n^-(q)$.

Suppose that n is even, $|\lambda_{z-1}(h)| = |\lambda_{z+1}(h)| = 0$ and h satisfies (i). Then h is conjugate to an element of only one of the groups $GO_n^+(q)$ and $GO_n^-(q)$.

Proof. See [18, p. 38, Case (C)(i,i')]. □

Next we establish a necessary and sufficient condition for h to lie in $\Gamma_n^\square(q)$ or $\Gamma_n^\boxtimes(q)$. Let $h \in GL_n(q)$, f be the characteristic polynomial of h , and suppose that $f = (z-1)^{n_1}(z+1)^{n_2}f_0$ with $(f_0, z^2-1) = 0$. Then h is conjugate to a block diagonal matrix

$$\text{diag}(h_1, h_{-1}, h_0) \tag{4.4}$$

with blocks of dimension n_1 , n_2 and $n - n_1 - n_2$, respectively, corresponding to this factorization of f . We refer to the matrix in (4.4) as a normal form of h .

Let $h \in \Gamma_n(q)$. We may replace h by its normal form $\text{diag}(h_1, h_{-1}, h_0)$ with blocks of dimensions n_1 , n_2 and $n - n_1 - n_2$, respectively, and denote by V_1 the subspace of F_q^n spanned by the first n_1 rows. Suppose that $h = gg^\tau$. Then $h^g = g^\tau g = h^\tau$. It follows that V_1g is h^τ -invariant and the characteristic polynomial of h^τ on V_1g is equal to $(z-1)^{n_1}$, and hence $V_1g = V_1$. The same is true for other blocks, and so g is also block diagonal with blocks g_1 , g_{-1} and g_0 of the same dimensions as h_1 , h_{-1} and h_0 , respectively (see also [8, Lemma 8.2]). Since $h_i = g_i g_i^\tau$ for $i = 1, -1, 0$ and $\det g = \det g_1 \det g_{-1} \det g_0$, it suffices to consider the following special cases: $f = (z-1)^n$, $f = (z+1)^n$, and $(f, z^2-1) = 1$.

Recall that for odd q , the group $\Omega_{2n}^\varepsilon(q)$ is the kernel of the spinor norm $\theta : SO_{2n}^\varepsilon(q) \rightarrow F_q^\times / (F_q^\times)^2$. The definition of the spinor norm in [16, pp. 163–165] implies the following way to calculate it (see also [2, Proposition 1.6.11]).

Lemma 4.4. *Let q be odd, n be even, $h \in SO_n^\varepsilon(q)$ and B be the matrix of the invariant symmetric bilinear form of $SO_n^\varepsilon(q)$. Suppose that $\det(E - h) \neq 0$. Then $\theta(h) \equiv \det((E - h)B) \pmod{(F_q^\times)^2}$.*

Lemma 4.5. *Let q be odd, n be even, $h \in \Gamma_n(q)$ and f be the characteristic polynomial of h .*

- (i) *If $f = (z + 1)^n$, then $h \notin \Gamma_n^\boxtimes(q)$.*
- (ii) *Let $f = (z - 1)^n$. If $\lambda_{z-1}(h)$ has no odd parts, then $h \notin \Gamma_n^\boxtimes(q)$; otherwise, $h \in \Gamma_n^\square(q) \cap \Gamma_n^\boxtimes(q)$.*
- (iii) *Let $(f, z^2 - 1) = 1$. Then h is conjugate to an element of $SO_n^\varepsilon(q)$ for some unambiguously defined ε , and $h \in \Gamma_n^\square(q)$ if and only if $-h$ is conjugate to an element of $\Omega_n^\varepsilon(q)$.*

Proof. Let $h = gg^\tau$. Possibilities for g in the first two cases are found in [8, Section 8], and we use this result for calculations. For brevity, we write $x \equiv y$ to denote that $x \equiv y \pmod{(F_q^\times)^2}$.

(i) The group $GL_n(q) \rtimes \langle \tau \rangle$ can be embedded into $GL_{2n}(q)$, and we consider the Jordan decomposition of $g\tau$ in the latter group. So $g\tau = ug_1\tau = g_1\tau u$, where u is unipotent and $(g_1\tau)^2 = -E$; i.e. g_1 is a skew-symmetric matrix. Since the determinant of a skew-symmetric matrix is a square and $\det u = 1$, we see that $\det g$ is also a square.

(ii) Similarly, $g\tau = ug_1\tau = g_1\tau u$, where u is unipotent and $(g_1\tau)^2 = E$; i.e. g_1 is a symmetric matrix. Furthermore, $h = u^2$ and the equality $ug_1\tau = g_1\tau u$ is equivalent to the condition that u preserves the bilinear form defining by g_1 . By Lemma 4.3, if $\lambda_{z-1}(u) = \lambda_{z-1}(h)$ has odd parts, then u is conjugate both to an element of $SO_n^+(q)$ and to an element of $SO_n^-(q)$, and so we can choose g_1 with any determinant, square or non-square. If $\lambda(u)$ has no odd parts, then u is conjugate to an element of $SO_n^+(q)$ only, and since the multiplicities of even parts are even, n is divisible by 4. In this case $\det g_1 \equiv \det E$.

(iii) The fact that h is conjugate to an element of $SO_n^\varepsilon(q)$, where ε is unambiguously defined, follows from Lemmas 4.1 and 4.3. By Lemma 4.4, we have $\theta(h) \equiv \det(E - h) \det B$, where B is the matrix of the symmetric bilinear form preserved by h . Observing that $\theta(-E) \equiv \det B$, we have

$$\theta(-h) \equiv \det(E - h) = \det(E - gg^\tau) = \det(g^\top - g) \det g^\tau \equiv \det g,$$

where the final equivalence holds because the determinant of the skew-symmetric matrix $g^\top - g$ is a square. The proof is complete. □

We are ready to find $\omega(\tau PGL_n(q))$ and $\omega(\tau PSL_n(q))$.

Lemma 4.6. *Let q and $n \geq 3$ be odd. Then*

$$\omega(\tau PSL_n(q)) = \omega(\tau PGL_n(q)) = 2 \cdot \omega(Sp_{n-1}(q)).$$

Proof. The first equality was established in (4.1). To prove the second one, we need to show that the set of the projective orders of elements of $\Gamma_n(q)$ is equal to $\omega(Sp_{n-1}(q))$.

Let $h \in \Gamma_n(q)$. Since n is odd, Lemma 4.1 implies that 1 is an eigenvalue of h . It follows that the projective order of h is equal to the ordinary order.

We show first that $|h| \in \omega(Sp_{n-1}(q))$. Let $\text{diag}(h_1, h_{-1}, h_0)$ be a normal form of h as in (4.4) and denote the dimension of h_1 by k . Then k is odd. By Lemmas 4.1 and 4.2, the

unipotent matrix h_1 is conjugate to an element of $SO_k(q)$ and the matrix $\text{diag}(h_{-1}, h_0)$ is conjugate to an element of $Sp_{n-k}(q)$. If $k = 1$, then there is nothing to prove. If $k > 1$, then $|h_1| \in \omega_p(SO_k(q)) = \omega_p(Sp_{k-1}(q))$, and hence $|h| \in \omega(Sp_k(q) \times Sp_{n-k}(q)) \subseteq \omega(Sp_{n-1}(q))$.

It follows from Lemmas 4.1 and 4.2 that a semisimple matrix lies in $\Gamma_n(q)$ if and only if it is conjugate to a matrix of the form $\text{diag}(1, h')$ with $h' \in Sp_{n-1}(q)$, and thus $\omega_{p'}(\Gamma_n(q)) = \omega_{p'}(Sp_{n-1}(q))$.

Let $a \in \omega(Sp_{n-1}(q))$ and $|a|_p = p^t > 1$. If $a = 2p^t$ and $n - 1 = p^{t-1} + 1$ (that is, the condition from (v) of Lemma 2.2 holds), then we define $h = \text{diag}(1, -J_{n-1})$. If $a = p^t$ or $n - 1 > p^{t-1} + 1$, then there is a semisimple matrix $h_s \in Sp_l(q)$, where $n - 1 = p^{t-1} + 1 + l$, such that $a = p^t|h_s|$, and we define $h = \text{diag}(J_{p^{t-1}+2}, h_s)$. It is easy to see that $h \in \Gamma_n(q)$ and $|h| = a$. The proof is complete. $\square$

Lemma 4.7. *Let q be odd and $n \geq 4$ be even. Then*

$$\begin{aligned} \omega(\tau PGL_n(q)) &= 2 \cdot \omega(PSp_n(q)), \\ \omega_{p'}(\tau PSL_n(q)) &= 2 \cdot \omega_{p'}(P\Omega_n^+(q)) \cup 2 \cdot \omega_{p'}(P\Omega_n^-(q)). \end{aligned}$$

If $n > 4$, then

$$\omega_{\bar{p}}(\tau PSL_n(q)) = 2 \cdot \omega_{\bar{p}}(\Omega_{n+1}(q)).$$

The set $\omega_{\bar{p}}(\tau PSL_4(q))$ consists of all multiples of p dividing $p(q \pm 1)$ if $p > 3$, and it consists of these multiples together with 18 if $p = 3$.

Proof. It suffices to find the sets of the projective orders of elements of $\Gamma_n(q)$ and $\Gamma_n^\square(q)$. For brevity, we denote these sets by $\omega(P\Gamma_n(q))$ and $\omega(P\Gamma_n^\square(q))$ respectively. Also we denote by Z the center of $GL_n(q)$.

Let $h \in \Gamma_n(q)$ and $\text{diag}(h_1, h_{-1}, h_0)$ be a normal form of h . Since n is even, it follows from Lemma 4.1 that the dimension of h_1 is even too. We denote this dimension by $2k$ and define $h_2 = \text{diag}(h_{-1}, h_0)$. Then h_1 is conjugate to an element of $SO_{2k}^+(q)$ and h_2 is conjugate to an element of $Sp_{n-2k}(q)$.

Let $k > 0$. Then $|h_1| \in \omega_p(SO_{2k}^+(q)) = \omega_p(Sp_{2k-2}(q))$, and hence

$$|h| \in \omega(Sp_{2k-2}(q) \times Sp_{n-2k}(q)) \subseteq \omega(Sp_{n-2}(q)).$$

Using Lemmas 2.2 and 2.3, it is easy to check that

$$\begin{aligned} \omega_{p'}(Sp_{n-2}(q)) &\subseteq \omega_{p'}(P\Omega_n^+(q)) \cup \omega_{p'}(P\Omega_n^-(q)) \subseteq \omega(PSp_n(q)), \\ \omega_{\bar{p}}(Sp_{n-2}(q)) &\subseteq \omega_{\bar{p}}(\Omega_{n+1}(q)) \subseteq \omega_{\bar{p}}(PSp_n(q)). \end{aligned}$$

Thus the projective order of h lies in the required sets.

Let $k = 0$. Then $h = h_2$ and $|hZ| \in \omega(PSp_n(q))$. It follows that

$$(P\Gamma_n(q)) \subseteq \omega(PSp_n(q)).$$

Suppose, in addition, that $h \in \Gamma_n^\square(q)$. Denote the dimension of h_{-1} by $2l$. By Lemma 4.5, we deduce that $h_{-1} \notin \Gamma_{2l}^\boxtimes(q)$, and so $-h_0$ is conjugate to an element of $\Omega_{n-2l}^\varepsilon(q)$ for some ε . Observe that $|hZ| = |-hZ|$.

Assume that $|-h_{-1}| > 1$. Then $|-hZ| = |-h| = |-h_{-1}| \cdot |-h_0|$. Since $|-h_{-1}| \in \omega_p(Sp_{2l}(q)) = \omega_p(\Omega_{2l+1}(q))$, we have

$$|hZ| \in \omega_{\bar{p}}(\Omega_{2l+1}(q) \times \Omega_{n-2l}^\varepsilon(q)) \subseteq \omega_{\bar{p}}(\Omega_{n+1}(q)).$$

Furthermore, if $n = 4$ then either $|hZ| = 9$ and $p = 3$, or $|hZ|$ lies in $p \cdot \Omega_2^\varepsilon(q)$ and, therefore, divides $p(q - \varepsilon)/2$.

If $|-h_{-1}| = 1$, then either $l = 0$ and $h = h_0$, or $l > 0$ and $|-hZ| = |-h_0|$. In either case, $|-hZ| \in \omega(P\Omega_n^\varepsilon(q)) \subseteq \omega(\Omega_{n+1}(q))$. In particular, if $n = 4$ and the order $|-hZ|$ is a multiple of p , then it divides $p(q \pm 1)/2$.

Now we prove the reverse containments. If h is semisimple, then $h \in \Gamma_n(q)$ if and only if h is conjugate to an element of $Sp_n(q)$, and hence

$$\omega_{p'}(PSp_n(q)) \subseteq \omega(P\Gamma_n(q)).$$

Let $a \in \omega(PSp_n(q))$ and $|a|_p = p^t > 1$. Lemma 2.2 implies that there is a semisimple matrix $h_s \in Sp_l(q)$, where $n = p^{t-1} + 1 + l$, such that $a = p^t|h_s|$, and we define $h = \text{diag}(-J_{p^{t-1}+1}, -h_s)$. It is easy to see that $h \in \Gamma_n(q)$ and $|hZ| = |-hZ| = |-h| = a$. Thus

$$\omega_{\bar{p}}(PSp_n(q)) \subseteq \omega(P\Gamma_n(q)),$$

and the assertion concerning $\tau PGL_n(q)$ follows.

Let $h \in \Omega_n^\varepsilon(q)$ be semisimple and $\text{diag}(h_1, h_{-1}, h_0)$ be a normal form of h , with h_1, h_{-1} of dimension k and l respectively. If $k > 0$, then $h \in \Gamma_n^\square(q)$ by Lemma 4.5(ii). If $k = 0$, then applying (i) or (iii) of Lemma 4.5 according as $l > 0$ or $l = 0$, we deduce that $-h \in \Gamma_n^\square(q)$. In any case, $|hZ| \in \omega(P\Gamma_n^\square(q))$.

Suppose that $n > 4$ and $a \in \omega_{\bar{p}}(\Omega_{n+1}(q))$, or $n = 4$ and $a = pc$ with c dividing $(q \pm 1)/2$, or $n = 4$, $p = 3$ and $a = 9$. Let $|a|_p = p^t$. By Lemmas 2.2 and 2.3, there are $\varepsilon \in \{+, -\}$ and a semisimple matrix $h_s \in \Omega_l^\varepsilon(q)$, where $n = p^{t-1} + 1 + l$, such that $a = p^t|h_s|$. Defining $h = \text{diag}(-J_{p^{t-1}+1}, -h_s)$, we see that $h \in \Gamma_n^\square(q)$ and $|hZ| = |-hZ| = |-h| = a$. The proof is complete. $\square$

In contrast to the sets of the projective orders of elements of $\Gamma_n(q)$ and $\Gamma_n^\square(q)$, the corresponding set for $\Gamma_n^\boxtimes(q)$, where n is even, is not in general closed under taking divisors. So we do not give an explicit description of the set $\omega(\tau\delta PSL_n(q))$ for even n . However, we derive some properties of this set.

Lemma 4.8. *Let q be odd and $n \geq 4$ be even. Then $\omega(\tau\delta PSL_n(q)) \not\subseteq \omega(PSL_n(q))$. If $k = 2^s$, where $s \geq 0$, and $q = q_0^k$, then $k \cdot \omega(\tau\delta(q_0)PSL_n(q_0)) \not\subseteq \omega(PSU_n(q))$. In particular, $\omega(\tau\delta(-q)PSU_n(q)) \not\subseteq \omega(PSL_n(q))$.*

Proof. Recall that $\omega(\tau\delta PSL_n(q))$ consists of the projective orders of matrices of $\Gamma_n^\boxtimes(q)$ multiplied by 2.

There is an element $h \in SO_n^\varepsilon(q) \setminus \Omega_n^\varepsilon(q)$ whose projective order is equal to $(q^{n/2} - \varepsilon)/2$. By Lemma 4.5, it follows that $-h \in \Gamma_n^\boxtimes(q)$, and so $q^{n/2} \pm 1 \in \omega(\tau\delta PSL_n(q))$.

Assume that $n > 4$ and $l = (n - 2)/2$ is odd. Then Lemma 2.2 together with the existence of primitive divisors $r_l(\pm q)$ implies that $p(q^l \pm 1) \in \omega(PSp_n(q)) \setminus \omega(\Omega_{n+1}(q))$. Applying Lemma 4.7, we see that

$$2p(q^l \pm 1) \in \omega(\tau PGL_n(q)) \setminus \omega(\tau PSL_n(q)) \subseteq \omega(\tau\delta PSL_n(q)). \quad (4.5)$$

Assume now that $n = 4$ and $q \equiv \varepsilon \pmod{4}$. The numbers $p(q + 1)$ and $p(q - \varepsilon)$ lie in $\omega(PSp_4(q))$ and do not divide $p(q \pm 1)/2$ (if $q - 1$ divides $(q + 1)/2$, then $q = 3$), and hence

$$2p(q + 1), 2p(q - \varepsilon) \in \omega(\tau\delta PSL_4(q)). \quad (4.6)$$

We prove the first assertion of the lemma and the second assertion for $k = 1$ together. Suppose that $\omega(\tau\delta PSL_n(q)) \subseteq \omega(PSL_n^\varepsilon(q))$ and as usual define $d = (n, q - \varepsilon)$. Then

$q^{n/2} + \varepsilon^{n/2} \in \omega(PSL_n^\varepsilon(q))$. By Lemma 2.7, this is equivalent to $(n)_2 > (q - \varepsilon)_2$. It follows that $l = (n - 2)/2$ is odd and $(d)_2 = (q - \varepsilon)_2$, and so $(q^l - \varepsilon)/d$ is odd. Then $2(q^l + \varepsilon)$ does not divide $(q^{n-2} - 1)/d$ and, therefore, $2p(q^l + \varepsilon) \notin \omega(PSL_n^\varepsilon(q))$. If $n > 4$, this contradicts (4.5). If $n = 4$, then $q \equiv -\varepsilon \pmod{4}$ and (4.6) implies that $2p(q + \varepsilon) \in \omega(\tau\delta PSL_4(q))$ yielding a contradiction.

Suppose now that $k > 1$ and $k \cdot \omega(\tau\delta(q_0)PSL_n(q_0)) \subseteq \omega(PSU_n(q))$. Observe that $(q + 1)_2 = 2$ and choose ϵ so that $q_0 \equiv \epsilon \pmod{4}$. Let $n/2$ be odd. By assumption, $a = k(q_0^{n/2} - \epsilon) \in \omega(PSU_n(q))$. Since a is a multiple of $r_{n/2}(\epsilon q_0)$ and $r_{n/2}(\epsilon q_0) \in R_{n/2}(q) = R_n(-q)$, it follows that a divides $c = (q^n - 1)/((q + 1)(n, q + 1))$. This is a contradiction because

$$(a)_2 = (q^{n/2} - 1)_2 > (c)_2 = (q^n - 1)_2/4.$$

Thus $l = (n - 2)/2$ is odd and by (4.5) and (4.6), we have $2p(q_0^l - \epsilon) \in \omega(\tau\delta PSL_n(q_0))$. Then $a = 2kp(q_0^l - \epsilon) \in \omega(PSU_n(q))$, and hence a divides $p(q^{n-2} - 1)/(n, q + 1)$. This contradicts $(a)_2 = (q^{n-2} - 1)_2$.

The final assertion follows from the second one and Lemma 3.3. □

We close this section with a proof of Theorem 1 and one of its corollaries.

Proof of Theorem 1. Recall that $L = PSL_n^\varepsilon(q)$, $d = (n, q - \varepsilon)$ and consider the difference $\omega(\tau L) \setminus \omega(L)$. We analyze separately two cases according as n is odd or even.

Let n be odd. By Lemma 4.6, the difference under consideration is equal to

$$2 \cdot \omega(Sp_{n-1}(q)) \setminus \omega(L).$$

We consider the numbers from 2.2 defining $\omega(Sp_{n-1}(q))$ in turn.

Let $a = 2p^t$, where $p^{t-1} + 1 = n - 1$. Any element of $\omega(L)$ that is a multiple of p^t divides $p^t(q - \varepsilon)/d$. Since d is odd, $2a$ divides $p^t(q - \varepsilon)/d$ if and only if 4 divides $q - \varepsilon$. Thus $4p^t \in \omega(\tau L) \setminus \omega(L)$ if and only if $n = p^{t-1} + 2$ for some $t \geq 1$ and $q \equiv -\varepsilon \pmod{4}$.

Let $a = p^t[q^{n_1} \pm 1, q^{n_2} \pm 1, \dots, q^{n_s} \pm 1]$, where $p^{t-1} + 1 + 2n_1 + 2n_2 + \dots + 2n_s = n - 1$, or $a = [q^{n_1} \pm 1, q^{n_2} \pm 1, \dots, q^{n_s} \pm 1]$, where $s \geq 2$ and $2(n_1 + n_2 + \dots + n_s) = n - 1$. Then $2a$ divides $p^t[q^{2n_1} - 1, q^{2n_2} - 1, \dots, q^{2n_s} - 1, q - \varepsilon]$ or $[q^{2n_1} - 1, q^{2n_2} - 1, \dots, q^{2n_s} - 1, q - \varepsilon]$ respectively.

The remaining possibility is $a = q^l \pm 1$, where $l = (n - 1)/2$. If $a = q^l + \varepsilon^l$, then $2a$ divides

$$\frac{q^{n-1} - 1}{d} = (q^l + \varepsilon^l) \frac{q^l - \varepsilon^l}{d}$$

since d is odd. Let $a = q^l - \varepsilon^l$. Assume that l is not a 2-power. Then $(l)_2 \leq l/3$, and hence $l + 2(l)_2 < n$. It follows that L has an element of order

$$c = [q^l - \varepsilon^l, q^{2(l)_2} - 1, q - \varepsilon].$$

Since $2(a)_2 = 2(q^l - \varepsilon^l)_2 \leq (q^{2l} - 1)_2 = (q^{2(l)_2} - 1)_2$, we have that $2a$ divides c . Finally, assume that l is a 2-power. Then the 2-exponent of L is equal to $(q^{2l} - 1)_2$ and any element of $\omega(L)$ that is a multiple of $(q^{2l} - 1)_2$ divides $(q^{2l} - 1)/d$. Clearly, $2a$ divides $(q^{2l} - 1)/d$ if and only if d divides $q^l + \varepsilon^l$, which is equivalent to $d = 1$. Thus $2(q^l - \varepsilon^l) \in \omega(\tau L) \setminus \omega(L)$ if and only if $n - 1 = 2^t$ and $d \neq 1$.

Let n be even. By Lemma 4.7, the set $\omega_p(\tau L)$ is equal to $2 \cdot \omega_p(\Omega_{n+1}(q))$ if $n > 4$, it consists of divisors of $p(q \pm 1)$ if $n = 4$ and $p > 3$, and it consists of divisors of $p(q \pm 1)$

together with 18 if $n = 4$ and $p = 3$. We can consider the numbers from Lemma 2.2 defining $\omega_{\bar{p}}(\Omega_{n+1}(q))$ for $n \geq 6$ and the numbers defining $\omega_{\bar{p}}(\tau L)$ for $n = 4$ together.

If $a = p^t$ and $n = p^{t-1} + 1$, then $2a \notin \omega(L)$.

Let $a = p^t[q^{n_1} \pm 1, q^{n_2} \pm 1, \dots, q^{n_s} \pm 1]$, where $s \geq 2$ and $p^{t-1} + 1 + 2n_1 + 2n_2 + \dots + 2n_s = n$. Then $2a$ divides $p^t[q^{2n_1} - 1, q^{2n_2} - 1, \dots, q^{2n_s} - 1] \in \omega(L)$.

Let $a = p^t(q^{n_1} - \epsilon)/2$, where $p^{t-1} + 1 + 2n_1 = n$. If $\epsilon = \epsilon^{n_1}$, then $2a$ divides $p^t(q^{n_1} - \epsilon^{n_1}) \in \omega(L)$, while if $\epsilon = -\epsilon^{n_1}$, then $2a$ divides

$$p^t \frac{q^{2n_1} - 1}{d} = p^t(q^{n_1} + \epsilon^{n_1}) \frac{q^{n_1} - \epsilon^{n_1}}{d}.$$

Thus

$$\omega_{\bar{p}}(\tau L) \subseteq \omega(L),$$

whenever $n \neq p^{t-1} + 1$.

To handle $\omega_{p'}(\tau L) = 2 \cdot \omega(P\Omega_{2n}^+(q)) \cup 2 \cdot \omega(P\Omega_{2n}^-(q))$, we consider the numbers from Lemma 2.3 that define $\omega_{p'}(P\Omega_{2n}^\pm(q))$.

Let $a = [q^{n_1} \pm 1, q^{n_2} \pm 1, \dots, q^{n_s} \pm 1]$, where $s \geq 3$ and $2(n_1 + n_2 + \dots + n_s) = n$. Then $2a$ divides $[q^{2n_1} - 1, q^{2n_2} - 1, \dots, q^{2n_s} - 1] \in \omega(L)$.

Let $a = (q^{n/2} - \epsilon)/(4, q^{n/2} - \epsilon)$. If $\epsilon = \epsilon^{n/2}$, then $2a$ divides $q^{n/2} - \epsilon^{n/2} \in \omega(L)$. If $\epsilon = -\epsilon^{n/2}$ and $(4, q^{n/2} + \epsilon^{n/2}) = 4$, then $2a$ divides

$$\frac{q^n - 1}{(q - \epsilon)d} = \frac{(q^{n/2} + \epsilon^{n/2})(q^{n/2} - \epsilon^{n/2})}{(q - \epsilon)d}$$

because $d/2$ divides $(n/2, q - \epsilon) = ((q^{n/2} - \epsilon^{n/2})/(q - \epsilon), q - \epsilon)$. If $\epsilon = -\epsilon^{n/2}$ and $(4, q^{n/2} + \epsilon^{n/2}) = 2$, then Lemma 2.7 implies that $2a = q^{n/2} + \epsilon^{n/2}$ does not lie in $\omega(L)$ if and only if $(n)_2 \leq (q - \epsilon)_2$. Assuming the last inequality, the condition $(4, q^{n/2} + \epsilon^{n/2}) = 2$ is equivalent to $q \equiv \epsilon \pmod{4}$.

Finally, let $a = [q^{n_1} - \epsilon_1, q^{n_2} - \epsilon_2]/e$, where $2(n_1 + n_2) = n$, $e = 2$ if $(q^{n_1} - \epsilon_1)_2 = (q^{n_2} - \epsilon_2)_2$ and $e = 1$ if $(q^{n_1} - \epsilon_1)_2 \neq (q^{n_2} - \epsilon_2)_2$. Define $n_0 = n/(2n_1, 2n_2)$. We may assume that $(q^{n_1} - \epsilon_1)_2 \geq (q^{n_2} - \epsilon_2)_2$.

If $\epsilon_2 = \epsilon^{n_2}$, then $2a$ divides $[q^{2n_1} - 1, q^{n_2} - \epsilon^{n_2}] \in \omega(L)$ since

$$2(a)_2 \leq 2(q^{n_1} - \epsilon_1)_2 \leq (q^{2n_1} - 1)_2.$$

Similarly, if $\epsilon_1 = \epsilon^{n_1}$ and $(q^{n_1} - \epsilon_1)_2 = (q^{n_2} - \epsilon_2)_2$, then $2a$ divides $[q^{2n_2} - 1, q^{n_1} - \epsilon^{n_1}]$.

Let $\epsilon_2 = -\epsilon^{n_2}$ and $\epsilon_1 = -\epsilon^{n_1}$. If $e = 2$, then $2a$ divides $[q^{2n_1} - 1, q^{2n_2} - 1]/(q - \epsilon)$. If $e = 1$, then n_1 and n_2 have opposite parity, therefore, $n/2$ is odd. It follows that n_0 is odd too, and $2a$ divides $[q^{2n_1} - 1, q^{2n_2} - 1]/(n_0, q - \epsilon)$.

Let $\epsilon_2 = -\epsilon^{n_2}$ and $\epsilon_1 = \epsilon^{n_1}$. We may assume that $(q^{n_1} - \epsilon^{n_1})_2 > (q^{n_2} + \epsilon^{n_2})_2$. If $n_1 \neq (n_1)_2$, then $n_1 + 2(n_1)_2 + 2n_2 < n$ and $2a$ divides $[q^{n_1} - \epsilon^{n_1}, q^{2n_2} - 1, q^{2(n_1)_2} - 1]$. If $2(q^{n_1} - \epsilon^{n_1})_2 \leq (q^{2n_2} - 1)_2$, then $2a$ divides $[q^{2n_2} - 1, q^{n_1} - \epsilon^{n_1}]$. If $q^{n_2} + \epsilon^{n_2}$ divides $q^{2n_1} - 1$, then $2a$ divides $q^{2n_1} - 1$ too. Finally, if $(n_0, q - \epsilon) = 1$, then $2a$ divides $[q^{2n_1} - 1, q^{2n_2} - 1]$.

Thus it remains to consider the case when $n_1 = (n_1)_2$, $2(q^{n_1} - \epsilon^{n_1})_2 > (q^{2n_2} - 1)_2$, $q^{n_2} + \epsilon^{n_2}$ does not divide $q^{2n_1} - 1$ and $(n_0, q - \epsilon) \neq 1$. The two first conditions yield $n_1 > (n_2)_2$. In particular, $(n/2)_2 = (n_2)_2$ and $n_0 = n/(2n_2)_2 = (n)_{2'}$. Then the third condition is equivalent to n_2 not being a 2-power. Observe that

$$(n)_{2'} = n/(2n_2)_2 = (n_1 + n_2)/(n_2)_2 = n_1/(n_2)_2 + (n_2)_{2'}.$$

It follows that $(n)_{2'}$ is a sum of a non-identity 2-power and an odd number greater than 1 and also $((n)_{2'}, q - \varepsilon) \neq 1$. In particular, $(n)_{2'} > 3$ and $(n, q - \varepsilon)_{2'} \neq 1$.

Conversely, suppose that $n = 2^t \cdot l$, where $t \geq 1$, $l \geq 5$ is odd and $(n, q - \varepsilon)$ is divisible by an odd prime r . Writing $n_1 = (n)_2 = 2^t$, $n_2 = n/2 - n_1 = 2^{t-1}(l - 2)$ and $a = [q^{n_1} - 1, q^{n_2} + \varepsilon^{n_2}]$, we see that

$$(q^{n_1} - 1)_2 = 2^{t-1}(q^2 - 1)_2 > (n_2)_2(q + \varepsilon)_2 \geq (q^{n_2} + \varepsilon^{n_2})_2$$

and hence $2a \in \omega(\tau L)$. Assume that $2a \in \omega(L)$ and let $c = [q^{l_1} - \varepsilon^{l_1}, \dots, q^{l_s} - \varepsilon^{l_s}]/f$, where $l_1 + \dots + l_s = n$, be a number from Lemma 2.1 that $2a$ divides. Since $2(a)_2 = 2(q^{n_1} - 1)_2 = (q^{2n_1} - 1)_2$, some of the numbers $l_1, \dots, l_s$ is a multiple of $2n_1$. Also $r_{2n_2}(\varepsilon q)$ divides a , and hence some of them is a multiple of $2n_2$. Observing that $[2n_1, 2n_2] = 2^{t+1}(l - 2) > 2^t l = n$, we deduce that those are different numbers, which yields $s = 2$, $l_1 = 2n_1$ and $l_2 = 2n_2$. Then $f = (n/(2n_1, 2n_2), q - \varepsilon)$, and so $(f)_r > 1$. Since $(r, q + \varepsilon) = (r, n_1) = (r, n_2) = 1$, it follows that

$$(c)_r = \frac{(q - \varepsilon)_r}{(f)_r} < (q - \varepsilon)_r = (a)_r.$$

This is a contradiction, therefore, $2a \notin \omega(L)$, and the proof is complete. $\square$

Theorem 1 has an interesting corollary: as the next lemma shows, if τ is admissible as an automorphism of $PSL_n^\varepsilon(q)$, then it is admissible as an automorphism of $PSL_n^\varepsilon(q^{1/k})$ for every odd k .

Lemma 4.9. *Let k be odd and $q = q_0^k$. If $\omega(\tau L) \subseteq \omega(L)$, then*

$$\omega(\tau PSL_n^\varepsilon(q_0)) \subseteq \omega(PSL_n^\varepsilon(q_0)).$$

In particular, if $\varepsilon = +$, τ and $\beta = \varphi^{m/k}$ are admissible for L , then $\beta\tau$ is also admissible.

Proof. Assume that $\omega(\tau PSL_n^\varepsilon(q_0)) \not\subseteq \omega(PSL_n^\varepsilon(q_0))$. This implies that the numbers n and q_0 satisfy the conditions of one of the items (i)–(v) of Theorem 1. Since k is odd, it follows that $(n, q_0 - \varepsilon)$ divides $(n, q - \varepsilon)$ and $(q - 1)_2 = (q_0 - 1)_2$, and thus the numbers n and q satisfy the same conditions. This contradicts the hypothesis that $\omega(\tau L) \subseteq \omega(L)$.

To prove the second assertion, it suffices to show that $\omega(\beta\tau L) \subseteq \omega(L)$. By Lemma 3.3, the proved containment and admissibility of β , we see that

$$\omega(\beta\tau L) = k \cdot \omega(\tau PSL_n^\varepsilon(q_0)) \subseteq k \cdot \omega(PSL_n^\varepsilon(q_0)) = \omega(\beta L) \subseteq \omega(L),$$

and the proof is complete. $\square$

5. ADMISSIBLE GROUPS

In this section, we will prove Theorems 2 and 3. As we mentioned, the theorems with $n = 3$ were proved in [19, 20], and so we assume that $n \geq 4$. Throughout the section, $q = p^m$ is odd, $L = PSL_n^\varepsilon(q)$ and G is a group such that $L < G \leq \text{Aut } L$. Also, we fix the numbers $d = (n, q - \varepsilon)$ and $b = ((q - \varepsilon)/d, m)_d$.

We begin with lemmas that holds for both linear and unitary groups. We say that a subgroup of $\text{Out } L$ is admissible if it is the image of an admissible group.

Lemma 5.1. *If $G \cap \text{Inndiag } L > L$, then $\omega(G) \neq \omega(L)$. In particular, admissible groups of $\text{Out } L$ are abelian and any non-trivial admissible subgroup of the group $\langle \bar{\delta} \rangle \rtimes \langle \bar{\tau} \rangle$ is conjugate in this group to $\langle \bar{\tau} \rangle$.*

Proof. By Lemma 2.1, if $|G \cap \text{Inndiag } L|/|L| = i > 1$, then G has an element of order $(q^n - \varepsilon^n)i/(q - \varepsilon)d$, which does not lie in $\omega(L)$. Thus admissible groups of $\text{Out } L$ can be embedded into the image of the group generated by φ and γ , which is abelian. The group $\langle \bar{\delta} \rangle \rtimes \langle \bar{\tau} \rangle$ is dihedral, and so every its subgroup that intersects trivially with $\langle \bar{\delta} \rangle$ is conjugate to $\langle \bar{\tau} \rangle$ or $\langle \bar{\delta}\bar{\tau} \rangle$, and in the latter case we may assume that n is even. But Lemma 4.8 says that $\langle \bar{\delta}\bar{\tau} \rangle$ is not admissible. $\square$

Lemma 5.2. *Suppose that $n \geq 5$ and $|G/L|$ is odd. Then $\omega(G) = \omega(L)$ if and only if $n - 1$ is not a p -power, G/L is conjugate in $\text{Out } L$ to a subgroup of $\langle \bar{\varphi} \rangle$ and $|G/L|$ divides $((q - \varepsilon)/d, m)_d$.*

Proof. If $\varepsilon = +$, the assertion is proved in [10, Propositions 6,7], and if $\varepsilon = -$, it is proved in [12, Proposition 6]. $\square$

Lemma 5.3. *Let $n = 4$ and $|G/L|$ is odd. Then $\omega(G) = \omega(L)$ if and only if 12 divides $q + \varepsilon$ and G/L is a 3-group.*

Proof. Let $\omega(G) = \omega(L)$ and $r \in \pi(G/L)$. Then G contains a field automorphism of order r , and hence by Lemma 3.3, $\omega(G)$ includes $r \cdot \omega(PSL_4^\varepsilon(q_0))$, where $q = q_0^r$. It follows that $rr_4(q_0), rr_3(\varepsilon q_0) \in \omega(L)$. Since $r_4(q_0) \in R_4(q)$, we see that r divides $a = (q^2 + 1)(q + \varepsilon)/d$ and, in particular, it does not divide $q - \varepsilon$. If $r \neq 3$, then $r_3(\varepsilon q_0) \in R_3(\varepsilon q)$, and so r divides $c = (q^3 - \varepsilon)/d$, which is a contradiction because $(a, c) = 1$. Let $r = 3$. Then 3 divides $q + \varepsilon$. Since $3p(q_0 - \varepsilon) \in \omega(L)$, we have that $3(q_0 - \varepsilon)$ divides $(q^2 - 1)/d$, and thus 4 divides $q + \varepsilon$.

Conversely, let 12 divide $q + \varepsilon$ and G/L be a 3-group. We may assume that G is the extension of L by a field automorphism of order dividing $(m)_3$. To prove that $\omega(G) \subseteq \omega(L)$, it suffices to check that $k \cdot \omega(PSL_4^\varepsilon(q_0)) \subseteq \omega(PSL_4^\varepsilon(q))$ with $q = q_0^k$ for every divisor k of $(m)_3$. Note that 12 divides $q_0 + \varepsilon$, $(q + \varepsilon)_3 = k(q_0 + \varepsilon)_3$ and $(4, q_0 - \varepsilon) = 2 = d$. Since $p \neq 3$, the set $\omega(PSL_4^\varepsilon(q_0))$ consists of divisors of $(q_0^2 + 1)(q_0 + \varepsilon)/2$, $(q_0^3 - \varepsilon)/2$, $q_0^2 - 1$ and $p(q_0^2 - 1)/2$. Observing that

$$k(q_0^2 + 1)(q_0 + \varepsilon) \mid (q^2 + 1)(q + \varepsilon),$$

$$k(q_0^3 - \varepsilon), k(q_0^2 - 1) \mid q^2 - 1,$$

we obtain the desired containment. $\square$

By Lemmas 5.2 and 5.3, it follows that up to conjugacy $\text{Out } L$ has only one maximal admissible subgroup of odd order, and we can take this subgroup to be $\langle \bar{\psi} \rangle$, where $\psi \in \langle \varphi \rangle$ has order $(b)_{2'}$ if $n > 4$ or $q \not\equiv 1 \pmod{12}$, and order $(m)_3$ if $n = 4$ and $q \equiv -\varepsilon \pmod{12}$.

Let $\eta = \delta^{(d)_{2'}}$ and S_2 be the Sylow 2-subgroup of $\text{Out } L$ generated by $\bar{\eta}$, $\bar{\varphi}^{(m)_{2'}}$ and $\bar{\tau}$.

Lemma 5.4. *If $\omega(G) = \omega(L)$, then G/L is conjugate in $\text{Out } L$ to a subgroup of $\langle \bar{\psi} \rangle \times S_2$.*

Proof. By Lemma 5.1, the group G/L is abelian, and hence is the direct product of its Hall $2'$ -subgroup A_1 and Sylow 2-subgroup A_2 . Since A_1 is admissible, it is conjugate to a subgroup of $\langle \bar{\psi} \rangle$. Replacing G/L by a conjugate if necessary, we may assume that $A_1 \leq \langle \bar{\psi} \rangle$.

Note that $\bar{\eta}$ centralizes $\bar{\psi}$. Indeed, if $\varepsilon = +$, then $\bar{\eta}^{\bar{\psi}} = \bar{\eta}^{q_0}$, where $q = q_0^{|\psi|}$. Since $|\psi|$ is odd, we have $(q - 1)_2 = (q_0 - 1)_2$, and hence $(d)_2$ divides $q_0 - 1$. If $\varepsilon = -$, then $\bar{\eta}^{\bar{\psi}} = \bar{\eta}^{q_0}$, where $q^2 = q_0^{|\psi|}$. Now $(q + 1)_2 < (q^2 - 1)_2 = (q_0 - 1)_2$, and again $(d)_2$ divides $q_0 - 1$. It

follows that the whole group S_2 centralizes $\overline{\psi}$. Thus A_2 is conjugate in $C_{\text{Out } L}(A_1)$ to a subgroup of S_2 , and the whole group G/L is conjugate to a subgroup of $\langle \overline{\psi} \rangle \times S_2$. $\square$

The structure of S_2 varies according to linear or unitary groups are under consideration, so in the rest of this section we consider the cases $\varepsilon = +$ and $\varepsilon = -$ separately. We begin with the case of unitary groups, in which $S_2 = \langle \overline{\eta} \rangle \rtimes \langle \overline{\varphi}^{(m)_{2'}} \rangle$.

Lemma 5.5. *Let $\varepsilon = -$ and $1 < G/L \leq S_2$. If $\omega(G) = \omega(L)$, then G/L is conjugate to a subgroup of $\langle \overline{\varphi}^{(m)_{2'}} \rangle$.*

Proof. We may assume that n is even. If m is odd, then $S_2 = \langle \overline{\eta} \rangle \rtimes \langle \overline{\tau} \rangle$ and by Lemma 5.1, the group G/L is conjugate to $\langle \overline{\tau} \rangle$.

Let m be even. Then $|\overline{\eta}| = 2$ and $S_2 = \langle \overline{\eta} \rangle \times \langle \overline{\varphi}^{(m)_{2'}} \rangle$. Suppose that $G \not\leq \langle \overline{\varphi}^{(m)_{2'}} \rangle$. Then G contains $\varphi^{m/k} \eta$ for some $k > 1$ dividing $(m)_2$. Let $q = q_0^k$. By Lemmas 3.3 and 4.8 together with Lemma 4.2, we have

$$\omega(\varphi^{m/k} \eta L) = k \cdot \omega(\tau \delta^{d/2}(q_0) PSL_n(q_0)) = k \cdot \omega(\tau \delta(q_0) PSL_n(q_0)) \not\subseteq \omega(L),$$

which is a contradiction. $\square$

Thus if $\varepsilon = -$ and τ is not admissible, then S_2 has no non-trivial admissible subgroups.

Lemma 5.6. *Suppose that $\varepsilon = -$, τ is admissible, $k > 1$ divides $(m)_2$ and $\beta = \varphi^{m/k}$. Then β is admissible if and only if $(n)_2 \leq 2$ or $n = 4, 8, 12$. If β is admissible and $\gamma \in \langle \varphi \rangle$ is admissible and has odd order, then $\gamma\beta$ is also admissible.*

Proof. We show first that $\omega(\beta L) \subseteq \omega(L)$ if and only if $(n)_2 \leq 2$ or $n = 4, 8, 12$. By Lemma 3.3, the set $\omega(\beta L)$ is equal to $k \cdot \omega(\tau PSL_n(q_0))$, where $q = q_0^k$.

Let n be odd. Since τ is admissible and $q \equiv 1 \pmod{4}$, Theorem 1 implies that $n - 2$ is not a power of p . Then using Lemmas 2.2 and 2.6, it is not hard to check that $k \cdot \omega(Sp_{n-1}(q_0)) \subseteq \omega(Sp_{n-1}(q))$. Now applying Lemma 4.6 yields

$$k \cdot \omega(\tau PSL_n(q_0)) = 2k \cdot \omega(Sp_{n-1}(q_0)) \subseteq 2 \cdot \omega(Sp_{n-1}(q)) = \omega(\tau L) \subseteq \omega(L).$$

Let n be even. Observe that $(d)_2 = (q + 1)_2 = 2$. By admissibility of τ and Theorem 1, it follows that $n - 1$ is not a p -power. Also it is not hard to verify that $k \cdot \omega(\Omega_{n+1}(q_0)) \subseteq \omega(\Omega_{n+1}(q))$ (cf. [11, Theorem 1]). Applying Lemma 4.7, we see that $k \cdot \omega_{\overline{p}}(\tau PSL_n(q_0)) \subseteq \omega_{\overline{p}}(\tau L)$ for $n > 4$. Since $k(q_0 \pm 1)$ divides $q - 1$, the same is true for $n = 4$. Thus it remains to examine when $2k \cdot \omega_{p'}(P\Omega_n^\pm(q_0)) \subseteq \omega(L)$, and we consider the numbers from Lemma 2.3 in turn.

Let $a = [q_0^{n_1} \pm 1, q_0^{n_2} \pm 1, \dots, q_0^{n_s} \pm 1]$, where $s \geq 3$ and $2(n_1 + n_2 + \dots + n_s) = n$. Then $2ka$ divides $[q^{2n_1} - 1, q^{2n_2} - 1, \dots, q^{2n_s} - 1] \in \omega(L)$.

Let $a = (q_0^{n/2} - \epsilon)/(4, q_0^{n/2} - \epsilon)$. If $n/2$ is even, then $2ka$ divides $q^{n/2} - 1 \in \omega(L)$. Suppose that $n/2$ is odd. Then $r_{n/2}(\epsilon q)$ divides a and lies in $R_{n/2}(q) = R_n(-q)$, and so $2ka \in \omega(L)$ id and only if $2ka$ divides

$$c = \frac{q^n - 1}{(q + 1)d} = \frac{(q^{n/2} - 1)(q^{n/2} + 1)}{(q + 1)d}.$$

Clearly, $(a)_{2'}$ divides $(c)_{2'}$. Since $(d)_2 = 2$, we see that $(c)_2 = (q - 1)_2/2$. If $(4, q_0 - \epsilon) = 4$, then $2k(a)_2 = k(q_0 - \epsilon)_2/2 = (q - 1)_2/2$. If $(4, q_0 - \epsilon) = 2$, then $2k(a)_2 = 2k \leq (q - 1)_2/2$.

Finally, let $a = [q_0^{n_1} - \epsilon_1, q_0^{n_2} - \epsilon_2]/e$, where $2(n_1 + n_2) = n$, $e = 2$ if $(q^{n_1} - \epsilon_1)_2 = (q^{n_2} - \epsilon_2)_2$ and $e = 1$ if $(q^{n_1} - \epsilon_1)_2 \neq (q^{n_2} - \epsilon_2)_2$.

Suppose that $n/2$ is odd. We may assume that $(q^{n_1} - \epsilon_1)_2 \geq (q^{n_2} - \epsilon_2)_2$. If n_2 is even, then $2ka$ divides $[q^{2n_1} - 1, q^{n_2} - 1]$. Let n_2 is odd. Then n_1 is even. If $n_1 \neq (n_1)_2$, then $2ka$ divides $[q^{n_1} - 1, q^{(n_1)_2} - 1, q^{2n_2} - 1]$. If $n_1 = (n_1)_2$ and $n_2 = 1$, then $2ka$ divides $q^{2n_1} - 1$. If $(n_1) = (n_1)_2$ and $n_2 > 1$, then by admissibility of τ , we have $(n, q+1)_{2'} = 1$ and, therefore, $(n/(2n_1, 2n_2), q+1) = 1$. So $[q^{2n_1} - 1, q^{2n_2} - 1]$ lies in $\omega(L)$, and it is divisible by $2ka$.

Suppose that $n/2$ is even and $n/2 \geq 8$. We can take n_1 and n_2 to be odd coprime numbers larger than 1. Also we take $\epsilon_1 = +$ and $\epsilon_2 = -$. Then $a = [q_0^{n_1} - 1, q_0^{n_2} + 1]$ and $2k(a)_2 = k(q_0^2 - 1)_2 = (q^2 - 1)_2$. Since a is a multiple of both $r_{n_1}(q_0)$ and $r_{n_2}(-q_0)$ and $r_{n_i}(\pm q_0) \in R_{2n_i}(-q)$ for $i = 1, 2$, it follows that $2ka \in \omega(L)$ if and only if $2ka$ divides $c = [q_{2n_1} - 1, q^{2n_2} - 1]/(n/(2n_1, 2n_2), q+1)$. But $(c)_2 = (q^2 - 1)_2/2 < (a)_2$ because $n/2$ is even. Thus $\omega(\beta L) \not\subseteq \omega(L)$.

We are left with the cases $n = 4, 8, 12$. If $n = 4$, then $2a$ divides $q_0^2 - 1$, and hence $2ka$ divides $q^2 - 1$. If $n = 8$, then a divides $[q_0^3 \pm 1, q_0 \pm 1]$ or $(q_0^4 - 1)/2$, and so $2ka$ divides $q^6 - 1$ or $q^4 - 1$. If $n = 12$, then a divides $[q_0^5 \pm 1, q_0 \pm 1]$, $[q_0^4 \pm 1, q_0^2 \pm 1]$, or $(q_0^6 - 1)/2$, and $2ka$ divides $q^{10} - 1$, $q^8 - 1$, or $q^6 - 1$.

We established that the condition for $\omega(\beta L)$ to be a subset of $\omega(L)$ depends only on n and not on q_0 , and thus β is admissible if and only if $\omega(\beta L) \subseteq \omega(L)$.

Next, suppose that β is admissible, $\gamma \in \langle \varphi \rangle$ is admissible and has odd order l and let $q_0 = q_1^l$. To prove that $\gamma\beta$ is admissible, it suffices to show that $\omega(\gamma\beta L) \subseteq \omega(L)$. By Lemma 4.9, if we regard τ as an automorphism of $PSU_n(q_1^k)$, it is still be admissible, and so by the above result, it follows that $k \cdot \omega(\tau PSL_n(q_1)) \subseteq \omega(PSU_n(q_1^k))$. By Lemma 3.3, we have

$$\omega(\gamma\beta L) = lk \cdot \omega(\tau PSL_n(q_1)) \subseteq l \cdot \omega(PSU_n(q_1^k)) = \omega(\gamma L) \subseteq \omega(L).$$

□

We are in position to prove Theorem Theorem 3.

Proof of Theorem 3. Suppose that $\omega(G) = \omega(L)$. By Lemmas 5.4 and 5.5, it follows that G/L is conjugate to a subgroup in $\langle \bar{\psi} \rangle \times \langle \bar{\varphi}^{(m)_{2'}} \rangle$. If $n-1$ is a p -power, then Lemmas 4.7, 5.2 and 5.3 implies that any non-trivial element of G/L is not admissible. If $n-1$ is not a p -power, then ψ is admissible, and applying Lemma 5.6 completes the proof.

□

Now we assume that $\varepsilon = +$, and so $S_2 = \langle \bar{\eta} \rangle \rtimes (\langle \bar{\varphi}^{m_{2'}} \rangle \times \langle \bar{\tau} \rangle)$. Given $1 \neq \beta \in \langle \varphi^{(m)_{2'}} \rangle$, we define β_ϵ with $\epsilon \in \{+, -\}$ by setting $\beta_+ = \beta$ and $\beta_- = \beta\tau$.

Lemma 5.7. *Let $\varepsilon = +$, $k > 1$ divides $(m)_2$ and $q = q_0^k$. Then for any $\epsilon \in \{+, -\}$, the following hold:*

- (i) *if n is odd or k does not divide $(q-1)/d$, then $k \cdot \omega(PSL_n^\epsilon(q_0)) \not\subseteq \omega(L)$;*
- (ii) *if n is even, k divides $(q-1)/d$ and $n = 1 + p^{t-1}$, then $k \cdot \omega(PSL_n^\epsilon(q_0)) \setminus \{kp^t\} \subseteq \omega(L)$;*
- (iii) *if n is even, k divides $(q-1)/d$ and $n-1$ is not a p -power, then $k \cdot \omega(PSL_n^\epsilon(q_0)) \subseteq \omega(L)$.*

In particular, if $\beta = \varphi^{m/k}$, then β_ϵ is admissible if and only if n is even, k divides $(q-1)/d$ and $n-1$ is not a p -power. If β_ϵ is admissible and $\gamma \in \langle \varphi \rangle$ is admissible and has odd order, then $\gamma\beta_\epsilon$ is also admissible.

Proof. If n is odd, then $2r_n(\epsilon q_0) \notin \omega(L)$ since $r_n(\epsilon q_0) \in R_n(q)$ and $(q^n - 1)/(q - 1)$ is odd. If n is even and k does not divide $(q - 1)/d$, then $kr_{n-1}(\epsilon q_0) \notin \omega(L)$. Indeed, otherwise the fact that $r_{n-1}(\epsilon q_0) \in R_{n-1}(q)$ implies that k divides $(q^{n-1} - 1)/d$. But $(q^{n-1} - 1)_2/(d)_2 = (q - 1)_2/(d)_2$, and (i) follows.

Let n be even and k divides $(q - 1)/d$. We claim that $ka \in \omega(L)$ for all $a \in \omega(PSL_n^\epsilon(q_0))$ except $a = p^t$ for $n = 1 + p^{t-1}$. The number a divides one of the numbers in items (i)–(v) of Lemma 2.1, and we consider these possibilities in turn.

If $a = (q_0^n - 1)/(q_0 - \epsilon)(n, q_0 - \epsilon)$, then ka divides $q^{n/2} - 1$. Similarly, if $a = [q_0^{n_1} - \epsilon^{n_1}, q_0^{n_2} - \epsilon^{n_2}]/(n/(n_1, n_2), q_0 - \epsilon)$, where $n_1 + n_2 = n$ and both n_1, n_2 are even, then ka divides $[q^{n_1/2} - 1, q^{n_2/2} - 1]$. If $a = [q^{n_1} - \epsilon^{n_1}, q^{n_2} - \epsilon^{n_2}]/(n/(n_1, n_2), q_0 - \epsilon)$, where n_1 and n_2 are odd, then ka divides $c = [q^{n_1} - 1, q^{n_2} - 1]/(n/(n_1, n_2), q - 1)$. Indeed, if $q_0 \equiv -\epsilon \pmod{4}$, then $k(a)_2 = k \leq (c)_2$, while if $q_0 \equiv \epsilon \pmod{4}$, then $k(q_0 - \epsilon)_2 = (q - 1)_2 \geq (k)_2(n)_2$, which implies that $(q_0 - \epsilon)_2 \geq (n)_2$ and $k(a)_2 = (q - 1)_2/(n)_2 = (c)_2$. Also, a divides c by Lemma 2.5. Finally, if $a = [q_0^{n_1} - \epsilon^{n_1}, \dots, q_0^{n_s} - \epsilon^{n_s}]$, where $s \geq 3$ and $n_1 + \dots + n_s = n$, or $a = p^t[q_0^{n_1} - \epsilon^{n_1}, \dots, q_0^{n_s} - \epsilon^{n_s}]$, where $s \geq 2$ and $1 + p^{t-1} + n_1 + \dots + n_s = n$, then ka divides $[q^{n_1} - 1, \dots, q^{n_s} - 1]$ or $p^t[q^{n_1} - 1, \dots, q^{n_s} - 1]$, respectively.

Let $\beta = \varphi^{m/k}$. By Lemma 3.3, we have $\omega(\beta_\epsilon L) = k \cdot \omega(PSL_n^\epsilon(q_0))$. Thus if β_ϵ is admissible, then n is even, k divides $(q - 1)/d$ and $n - 1$ is not a p -power. Conversely, if all these three conditions are satisfied, then by the above

$$\omega(\beta^{k_1} L) = (k/k_1) \cdot \omega(PSL_n(q_0^{k_1})) \subseteq \omega(L)$$

for every divisor k_1 of k , and hence β_ϵ is admissible.

To prove the final assertion, it suffices to check that $\omega(\gamma\beta_\epsilon L) \subseteq \omega(L)$ for admissible γ and β_ϵ . Let $|\gamma| = l$ and $q_0 = q_1^l$. Observing that $(q - 1)_2 = (q_1^k - 1)_2$, we see that k divides $(q - 1)/d$ if and only if it divides $(q_1^k - 1)/(n, q_1^k - 1)$. Thus admissibility of β_ϵ yields $k \cdot \omega(PSL_n^\epsilon(q_1)) \subseteq \omega(PSL_n(q_1^k))$. Applying Lemma 3.3, we have

$$\omega(\gamma\beta_\epsilon L) = lk \cdot \omega(PSL_n^\epsilon(q_1)) \subseteq l \cdot \omega(PSL_n(q_1^k)) = \omega(\gamma L) \subseteq \omega(L),$$

and the proof is complete. $\square$

Lemma 5.8. *Let $\epsilon = +$, n be even, $k > 1$ divide $(m)_2$, $\beta = \varphi^{m/k}$ and $q = q_0^k$. If $\alpha = \beta_\epsilon \eta^j$ is admissible, then $\bar{\alpha}$ is conjugate in S_2 to either $\bar{\beta}_\epsilon$ or $\bar{\beta}_\epsilon \bar{\eta}$, with $q_0 \equiv -\epsilon \pmod{4}$ and $2k \leq (q - 1)_2/(d)_2$ in the latter case. Furthermore, if $q_0 \equiv -\epsilon \pmod{4}$ and $2k \leq (q - 1)_2/(d)_2$, then*

- (i) $\beta_\epsilon \eta$ is admissible if and only if n cannot be represented as $1 + p^{t-1} + 2^u$ with $t, u \geq 1$ and, in addition, $k = 2$ whenever $n - 1$ is a p -power;
- (ii) if $\beta_\epsilon \eta$ is admissible and $\gamma \in \langle \varphi \rangle$ is admissible and has odd order, then $\gamma\beta_\epsilon \eta$ is admissible.

Proof. Note that $\bar{\eta}^{\beta_\epsilon} = \bar{\eta}^{\epsilon q_0}$, and so

$$\bar{\beta}_\epsilon^{\bar{\eta}} = \bar{\eta}^{-1} \bar{\beta}_\epsilon \bar{\eta} = \bar{\beta}_\epsilon \bar{\eta}^{1 - \epsilon q_0}. \quad (5.1)$$

Suppose that $\alpha = \beta_\epsilon \eta^j$ is admissible and $\bar{\alpha}$ is not conjugate in S_2 to $\bar{\beta}_\epsilon$. Then (5.1) implies that $(q_0 - \epsilon, (d)_2)$ does not divide j , or in other words

$$(j)_2 < (q_0 - \epsilon, d)_2. \quad (5.2)$$

Since $\eta = \delta^{(d)_{2'}}$, applying Lemma 3.3 yields

$$\omega(\alpha L) = k \cdot \omega(\delta^{j(d)_{2'}}(\epsilon q_0) PSL_n^\epsilon(q_0)). \quad (5.3)$$

The index of $\delta^{j(d)_{2'}}(\epsilon q_0)PSL_n^\epsilon(q_0)$ in $PGL_n^\epsilon(q_0)$ is equal to $(j(d)_{2'}, q_0 - \epsilon)$. There is an element of order $q_0^{n-1} - \epsilon$ in $PGL_n(q_0)$, and hence writing $a = (q_0^{n-1} - \epsilon)/(j(d)_{2'}, q_0 - \epsilon)$, we see that $ka \in \omega(\alpha L)$. Since a is a multiple of $r_{n-1}(\epsilon q_0) \in R_{n-1}(q)$ and lies in $\omega(L)$, it follows that a divides $c = (q^{n-1} - 1)/d$. If $q_0 \equiv \epsilon \pmod{4}$, then by (5.2), we deduce that

$$(a)_2 = k(q_0 - \epsilon)_2/(j)_2 = (q - 1)_2/(j)_2 > (q - 1)_2/(d)_2 = (c)_2.$$

Thus $q_0 \equiv -\epsilon \pmod{4}$. Then $(q_0 - \epsilon, d)_2 = 2$, and so j is odd. In particular, (5.1) shows that α is conjugate to $\beta_\epsilon \eta$. Also, $(a)_2 = 2k \leq (c)_2 = (q - 1)_2/(d)_2$, and the first assertion is proved.

Next, let $\alpha = \beta_\epsilon \eta$, $q_0 \equiv -\epsilon \pmod{4}$ and $2k \leq (q - 1)_2/(d)_2$. Observing that $(q - 1)_2 = k(q_0 + \epsilon)_2$, we deduce from the last inequality that $2(d)_2 \leq (q_0 + \epsilon)_2$, and hence $\bar{\alpha}^2 = \bar{\beta}^2 \bar{\eta}^{1+\epsilon q_0} = \bar{\beta}^2$. Since $2k \leq (q - 1)_2/(d)_2$, it follows from Lemma 5.7 that β^2 is admissible if and only if $k = 2$ or $n - 1$ is not a p -power.

Suppose that $n = 1 + p^{t-1} + 2^u$ with $t, u \geq 1$. Then $PGL_n^\epsilon(q_0)$ has an element of order $p^t(q_0^{2^u} - 1)$, and so $ka \in \omega(\alpha L)$, where $a = p^t(q_0^{2^u} - 1)/(d, q_0 - \epsilon)_{2'}$. Since ka is a multiple of $k p^t(q_0^{2^u} - 1)_2 = p^t(q^{2^u} - 1)_2$, it lies in $\omega(L)$ only if it divides $p^t(q^{2^u} - 1)/d$, which is not the case.

To prove (i), it remains to verify that $\omega(\alpha L) \subseteq \omega(L)$ whenever n cannot be represented as $1 + p^{t-1} + 2^u$ with $t, u \geq 1$. Since $((d)_{2'}, q_0 - \epsilon) = (n, q_0 - \epsilon)_{2'} = (n, q_0 - \epsilon)/2$, it follows from (5.3) that

$$\omega(\alpha L) = k \cdot \omega(2PSL_n^\epsilon(q_0) \setminus PSL_n^\epsilon(q_0)),$$

where $2PSL_n^\epsilon(q_0)$ denotes the unique subgroup of $PGL_n^\epsilon(q_0)$ that contains $PSL_n(q_0)$ as a subgroup of index 2. Denote $\omega(2PSL_n^\epsilon(q_0) \setminus PSL_n^\epsilon(q_0))$ by ω .

If $n = p^{t-1} + 1$, then any element of $GL_n(q)$ whose order is divisible by p^t is conjugate to a scalar multiple of the $n \times n$ unipotent Jordan block, therefore, $p^t \notin \omega$. By Lemma 5.7, we have

$$k \cdot (\omega \cap \omega(PSL_n^\epsilon(q_0))) \subseteq \omega(L),$$

and hence we are left with elements of $\omega \setminus \omega(PSL_n^\epsilon(q_0))$. By Lemma 2.1, it follows that this difference consists of some divisors of the following numbers:

$$\begin{aligned} & 2(q_0^n - 1)/((q_0 - \epsilon)(n, q_0 - \epsilon)), \\ & [q_0^{n_1} - \epsilon^{n_1}, q_0^{n_2} - \epsilon^{n_2}]/(n/(n_1, n_2), q_0 - \epsilon), \text{ where } n_1, n_2 \text{ are odd and } n_1 + n_2 = n; \\ & 2p^t(q_0^{n_1} - \epsilon^{n_1})/(n, q_0 - \epsilon), \text{ where } n_1 > 0 \text{ and } p^{t-1} + 1 + n_1 = n. \end{aligned}$$

Let $a = 2(q_0^n - 1)/((q_0 - \epsilon)(n, q_0 - \epsilon))$. Then

$$k(a)_2 = 2k \frac{(q_0^n - 1)_2}{(q_0 - \epsilon)_2(q_0 - \epsilon, n)_2} = (q^n - 1)_2/2 = (q^{n/2} - 1)_2,$$

and so ka divides $q^{n/2} - 1 \in \omega(L)$.

Let $a = 2[q_0^{n_1} - \epsilon^{n_1}, q_0^{n_2} - \epsilon^{n_2}]/(n/(n_1, n_2), q_0 - \epsilon)$, where n_1, n_2 are odd and $n_1 + n_2 = n$. Consider the number $c = [q^{n_1} - 1, q^{n_2} - 1]/(n/(n_1, n_2), q - 1)$ lying in $\omega(L)$. By Lemma 2.5, both numbers $(q^{n_i} - 1)/(n/(n_1, n_2), q_0 - \epsilon)$, where $i = 1, 2$, divide c . Also, $k(a)_2 = 2k \leq (q - 1)_2/d_2 = (c)_2$, and hence ka divides c .

Let $a = 2p^t(q_0^{n_1} - \epsilon^{n_1})/(n, q_0 - \epsilon)$, where $n_1 > 0$ and $p^{t-1} + 1 + n_1 = n$. Note that n_1 is even and by hypothesis, $(n_1)_2 \neq n_1$. Since

$$k(a)_2 = 2k(q_0^{n_1} - 1)_2/(q_0 - \epsilon)_2 = (q^{n_1} - 1)_2 = (q^{(n_1)_2} - 1)_2,$$

we see that ka divides $p^t[q^{n_1}/2 - 1, q^{(n_1)_2} - 1] \in \omega(L)$.

To prove (ii), take admissible γ and α . We may assume that $n - 1$ is not a p -power because otherwise $\gamma = 1$ by Lemmas 5.2 and 5.3. Then by the equality $\bar{\alpha}^2 = \bar{\beta}^2$ and Lemma 5.8, to prove that $\gamma\alpha$ is admissible, it suffices to check that $\omega(\gamma\alpha L) \subseteq \omega(L)$. Denote $|\gamma|$ by l and let $q_0 = q_1^l$. Observe that $q_1 \equiv q_0 \equiv -\epsilon \pmod{4}$ and $2k$ divides $(q_1^k - 1)_2 / (n, q_1^k - 1)_2$.

By Lemma 3.3, we have

$$\omega(\gamma\alpha L) = lk \cdot \omega(\delta^{(d)_{2'}}(\epsilon q_1) PSL_n^\epsilon(q_1)).$$

Using the fact that $(d, q_1 - \epsilon)_{2'} = (q_1 - \epsilon)/2$ and the above results, we deduce that

$$k \cdot \omega(\delta^{(d)_{2'}}(\epsilon q_1) PSL_n^\epsilon(q_1)) = k \cdot \omega(2PSL_n^\epsilon(q_1) \setminus PSL_n^\epsilon(q_1)) \subseteq \omega(PSL_n(q_1^k)).$$

Thus

$$\omega(\gamma\alpha L) \subseteq l \cdot \omega(PSL_n(q_1^k)) = \omega(\gamma L) \subseteq \omega(L),$$

and the proof is complete. $\square$

We are now ready to describe admissible 2-subgroups of $\text{Out } L$ in the case of linear groups, and then prove Theorem 2. Recall that $b = ((q - \epsilon)/d, m)_d$ and define $\phi = \varphi^{m/(b)_2}$.

Lemma 5.9. *Let $\epsilon = +$ and $1 < G/L \leq S_2 = \langle \bar{\eta} \rangle \rtimes (\langle \bar{\varphi}^{(m)_{2'}} \rangle \times \langle \bar{\tau} \rangle)$. Then $\omega(G) = \omega(L)$ if and only if G/L is cyclic and up to conjugacy in S_2 is generated by the image of one of the following elements:*

- (i) τ if b is odd and τ is admissible;
- (ii) $\beta_\pm$, where $1 \neq \beta \in \langle \phi \rangle$, if b is even and $n - 1$ is not a p -power;
- (iii) $\beta\tau\eta$, where $1 \neq \beta \in \langle \phi^2 \rangle$, if $(b)_2 > 2$ and n cannot be represented as $1 + p^{t-1}$ or $1 + p^{t-1} + 2^u$ with $t, u \geq 1$;
- (iv) $\phi_{-\kappa}\eta$, where $p \equiv \kappa \pmod{4}$, if $(p - \kappa)_2 > (n)_2$ and n cannot be represented as $1 + p^{t-1}$ or $1 + p^{t-1} + 2^u$ with $t, u \geq 1$;
- (v) $\varphi^{m/2}\tau\eta$, if $(b)_2 > 2$, $n = 1 + p^{t-1}$ for some $t \geq 1$ and n cannot be represented as $2 + 2^u$ with $u \geq 1$;
- (vi) $(\varphi^{m/2})_{-\kappa}\eta$, where $p \equiv \kappa \pmod{4}$, if $(m)_2 = 2$, $(p - \kappa)_2 > (n)_2$, $n = 1 + p^{t-1}$ for some $t \geq 1$ and n cannot be represented as $2 + 2^u$ with $u \geq 1$.

Proof. Observe that b is even if and only if all the numbers n , m and $(q - 1)/d$ are even.

Suppose that n is odd. Then $S_2 = \langle \bar{\varphi}^{(m)_{2'}} \rangle \times \langle \bar{\tau} \rangle$. By Lemma 5.7, any non-trivial element of $\langle \varphi^{(m)_{2'}} \rangle$ is not admissible. and thus the only potentially admissible subgroup of S_2 is $\langle \bar{\tau} \rangle$.

Suppose that n is even but at least one of m and $(q - 1)/d$ is odd. It follows from Lemmas 5.7 and 5.8 that every admissible subgroup of S_2 is contained in $\langle \bar{\eta} \rangle \rtimes \langle \bar{\tau} \rangle$. Then by Lemma 5.1, it is conjugate to a subgroup of $\langle \bar{\tau} \rangle$, and we are done in the case of odd b .

Suppose now that all the numbers n , m and $(q - 1)/d$ are even. In particular, $(q - 1)_2 > (d)_2 = (n)_2$ and so τ is not admissible by Theorem 4.7. Then by Lemma 5.1, we obtain that all non-trivial elements of $\langle \bar{\eta} \rangle \rtimes \langle \bar{\tau} \rangle$ are not admissible. It follows that every admissible subgroup of S_2 is isomorphic to a subgroup of $\langle \bar{\varphi}^{(m)_{2'}} \rangle$, and hence is cyclic.

Let $\bar{\alpha} \in S_2 \setminus \langle \bar{\eta} \rangle \rtimes \langle \bar{\tau} \rangle$ is admissible. We may assume that $\alpha = \beta_\epsilon \eta^j$, where $\beta = \varphi^{m/k}$ for some $1 < k = 2^l \leq (m)_2$. By Lemmas 5.7 and 5.8, admissibility of α yields $k \leq (q - 1)_2 / (d)_2$ or, equivalently, $\beta \in \langle \phi \rangle$. Moreover, all conjugates of $\bar{\beta}_\epsilon$ are admissible whenever $n - 1$ is not a p -power, and this gives the automorphisms in (ii).

It remains to consider the case when $\bar{\alpha}$ is not conjugate to $\bar{\beta}_\epsilon$. In this case by Lemma 5.8, we may assume that $\alpha = \beta_\epsilon \eta$,

$$2k \leq (q-1)_2/(d)_2 \quad (5.4)$$

and $q_0 \equiv -\epsilon \pmod{4}$, where $q = q_0^k$. Moreover, $k = 2$ whenever $n-1$ is a p -power. Note that $(q-1)_2 = (m)_2(p-\kappa)_2$, where $p \equiv \kappa \pmod{4}$.

Suppose that $(q-1)_2/(d)_2 \leq (m)_2$ or, in other words, $(p-\kappa)_2 \leq (n)_2$. Then $k \leq (m)_2/2$, so $q_0 \equiv 1 \pmod{4}$ and the condition (5.4) is equivalent to $\beta \in \langle \phi^2 \rangle$. Thus $(b)_2 > 2$ and $\alpha = \beta\tau\eta$ with $\beta \in \langle \phi^2 \rangle$. In particular, if $n-1$ is a p -power, then $\alpha = \varphi^{m/2}\tau\eta$.

Suppose that $(q-1)_2/(d)_2 > (m)_2$ or, equivalently, $(p-\kappa)_2 > (n)_2$. Then the condition (5.4) holds for all $\beta \in \langle \phi \rangle$. Also, $q_0 \equiv 1 \pmod{4}$ if $k < (m)_2$ and $q_0 \equiv \kappa \pmod{4}$ if $k = (m)_2$. Thus $\alpha = \beta\tau\eta$ with $\beta \in \langle \phi^2 \rangle$ or $\alpha = \phi_{-\kappa}\eta$. In particular, if $n-1$ is a p -power, then $\alpha = \varphi^{m/2}\tau\eta$ when $(b)_2 = (m)_2 > 2$ and $\alpha = \phi_{-\kappa}\eta$ when $(b)_2 = (m)_2 = 2$.

Conversely, let α be one of the resulting automorphisms and let the associated conditions be satisfied. Then by Lemma 5.8, α is admissible unless n cannot be represented as $1 + p^{t-1} + 2^u$. If $n = 1 + p^{t-1}$ and $n = 1 + p^{s-1} + 2^u$, then $p^{t-1} = p^{s-1} + 2^u$ or, equivalently, $p^{s-1}(p^{t-s} - 1) = 2^u$, and so $s = 1$ and $n = 2 + 2^u$. Thus we obtain the automorphisms in (iii)–(vi), and the proof is complete. $\square$

Proof of Theorem 2. Let $\omega(G) = \omega(L)$. By Lemma 5.4, it follows that G/L is conjugate to a subgroup of $\langle \bar{\psi} \rangle \times S_2$. Admissible subgroups of $\langle \bar{\psi} \rangle$ and S_2 are described in Lemmas 5.2, 5.3 and Lemma 5.9 respectively. These descriptions together with Lemmas 5.7 and 5.8 imply that the product of an admissible subgroup of $\langle \bar{\psi} \rangle$ and an admissible subgroup of S_2 is also admissible, and this completes the proof. $\square$

REFERENCES

- [1] R. Brandl and W.J. Shi, The characterization of $PSL(2, q)$ by its element orders, *J. Algebra* **163** (1994), no. 1, 109–114.
- [2] J. Bray, D. Holt, and C. Roney-Dougall, The maximal subgroups of the low-dimensional finite classical groups, London Mathematical Society Lecture Note Series, vol. 407, Cambridge University Press, Cambridge, 2013.
- [3] A. A. Buturlakin, Spectra of finite linear and unitary groups, *Algebra Logic* **47** (2008), no. 2, 91–99.
- [4] A. A. Buturlakin, Spectra of finite symplectic and orthogonal groups, *Siberian Adv. Math.* **21** (2011), no. 3, 176–210.
- [5] A. A. Buturlakin and M. A. Grechkoseeva, The cyclic structure of maximal tori of the finite classical groups, *Algebra Logic* **46** (2007), no. 2, 73–89.
- [6] J. H. Conway, R. T. Curtis, S. P. Norton, R. A. Parker, and R. A. Wilson, Atlas of finite groups, Clarendon Press, Oxford, 1985.
- [7] J. Fulman and R. Guralnick, Bounds on the number and sizes of conjugacy classes in finite Chevalley groups with applications to derangements, *Trans. Amer. Math. Soc.* **364** (2012), no. 6, 3023–3070.
- [8] Jason Fulman and Robert Guralnick, Conjugacy class properties of the extension of $GL(n, q)$ generated by the inverse transpose involution, *J. Algebra* **275** (2004), no. 1, 356–396.
- [9] Rod Gow and C. Ryan Vinroot, Extending real-valued characters of finite general linear and unitary groups on elements related to regular unipotents, *J. Group Theory* **11** (2008), no. 3, 299–331.
- [10] M. A. Grechkoseeva, Recognition by spectrum for finite linear groups over fields of characteristic 2, *Algebra Logic* **47** (2008), no. 4, 229–241.
- [11] M. A. Grechkoseeva, On spectra of almost simple groups with symplectic or orthogonal socle, *Siberian Math. J.* **57** (2016), no. 4, 582–588.
- [12] M. A. Grechkoseeva and W. J. Shi, On finite groups isospectral to finite simple unitary groups over fields of characteristic 2, *Sib. Élektron. Mat. Izv.* **10** (2013), 31–37.

- [13] M. A. Grechkoseeva and A. V. Vasil'ev, On the structure of finite groups isospectral to finite simple groups, *J. Group Theory* **18** (2015), no. 5, 741–759.
- [14] P. Kleidman and M. Liebeck, The subgroup structure of the finite classical groups, London Mathematical Society Lecture Note Series, vol. 129, Cambridge University Press, Cambridge, 1990.
- [15] V. D. Mazurov, On the set of orders of elements of a finite group, *Algebra and Logic* **33** (1994), no. 1, 49–55.
- [16] D. E. Taylor, The geometry of the classical groups, Sigma Series in Pure Mathematics, vol. 9, Heldermann Verlag, Lemgo, 2009.
- [17] A. V. Vasil'ev, On finite groups isospectral to simple classical groups, *J. Algebra* **423** (2015), 318–374.
- [18] G. E. Wall, On the conjugacy classes in the unitary, symplectic and orthogonal groups, *J. Aust. Math. Soc.* **3** (1963), 1–62.
- [19] A. V. Zavarnitsine, Recognition of the simple groups $L_3(q)$ by element orders, *J. Group Theory* **7** (2004), no. 1, 81–97.
- [20] A. V. Zavarnitsine, Recognition of the simple groups $U_3(q)$ by element orders, *Algebra Logic* **45** (2006), no. 2, 106–116.
- [21] K. Zsigmondy, Zur Theorie der Potenzreste, *Monatsh. Math. Phys.* **3** (1892), 265–284.
- [22] M. A. Zvezdina, On spectra of automorphic extensions of finite simple exceptional groups of Lie type, to appear in *Algebra Logic*.
- [23] M. A. Zvezdina, Spectra of automorphic extensions of finite simple symplectic and orthogonal groups over fields of characteristic 2, *Sib. Élektron. Mat. Izv.* **11** (2014), 823–832.

SOBOLEV INSTITUTE OF MATHEMATICS, AC. KOPTYUGA 4, NOVOSIBIRSK, 630090, RUSSIA; NOVOSIBIRSK STATE UNIVERSITY, PIROGOVA 4, NOVOSIBIRSK, 630090, RUSSIA

E-mail address: grechkoseeva@gmail.com